

7-15-2006

10,10'-Dinitro-10,10'-(butane-1,4-diyl)dianthracen-9(10H)-one

Mustafa Arslan

Sakarya University, Adapazari, Turkey

Erol Asker

Balikesir University, Balikesir, Turkey

John Masnovi

Cleveland State University, j.masnovi@csuohio.edu

Ronald J. Baker

Follow this and additional works at: http://engagedscholarship.csuohio.edu/scichem_facpub

 Part of the [Organic Chemistry Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

Asker, E., Masnovi, J., Baker, R. J., & Arslan, M. (2006). 10,10'-dinitro-10,10'-(butane-1,4-diyl)dianthracen-9(10H)-one. *Acta Crystallographica Section E*, 62(7), o2819-o2821. doi:10.1107/S160053680602157X

This Article is brought to you for free and open access by the Chemistry Department at EngagedScholarship@CSU. It has been accepted for inclusion in Chemistry Faculty Publications by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.

**10,10'-Dinitro-10,10'-(butane-1,4-diyl)-
dianthracen-9(10*H*)-one****Mustafa Arslan,^{a*} Erol Asker,^b
John Masnovi^c and
Ronald J. Baker^c**^aDepartment of Chemistry, Faculty of Arts and Sciences, Sakarya University, 54140 Esentepe/Adapazari, Turkey, ^bNecatibey Faculty of Education, Balikesir University, 10100 Balikesir, Turkey, and ^cDepartment of Chemistry, Cleveland State University, Cleveland, OH 44115, USA


Correspondence e-mail: marslan@sakarya.edu.tr

Key indicatorsSingle-crystal X-ray study
T = 295 K
Mean $\sigma(\text{C}-\text{C}) = 0.003 \text{ \AA}$
R factor = 0.042
wR factor = 0.110
Data-to-parameter ratio = 12.7For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

The title compound, $\text{C}_{32}\text{H}_{24}\text{N}_2\text{O}_6$, was obtained as the decomposition product of (*E,E*)-1,4-bis[9,10-dihydro-9-nitro-10-(trinitromethyl)-9-anthryl]butane, which was synthesized via a photochemical reaction of 1,4-bis(9-anthryl)butane with tetranitromethane. The asymmetric unit contains one half-molecule; the complete molecule is generated by a center of inversion. The crystal packing is determined mainly by intermolecular C—H...O interactions.

Received 13 April 2006
Accepted 7 June 2006**Comment**

Photonitration of aromatic compounds through the use of tetranitromethane (TNM) offers an alternative route to the conventional nitration processes which require the use of concentrated nitric and sulfuric acids. (Kochi, 1991; Butts *et al.*, 1996; Cox, 1998; Lehnig & Schürmann, 1998). In general, 9-alkyl substituted anthracene compounds lead to addition of a nitro group at the C-atom bearing an alkyl group and trinitromethylation takes place at the sterically less hindered unsubstituted C10 center. Photolysis of the 1, 4-bis(9-anthryl)butane/TNM charge transfer complex led to unstable (*E,E*)-1,4-bis[9,10-dihydro-9-nitro-10-(trinitromethyl)-9-anthryl]butane. When this was passed through a column of basic alumina or silica gel, trinitromethyl groups were eliminated to give the corresponding anthrone derivative. In this paper we report the crystal structure of the title compound, (I), as the decomposition product of this process.



The asymmetric unit contains one half-molecule; the complete molecule is generated by a center of inversion. Bond lengths and angles in the anthracene system (Table 1) are in agreement with those of related compounds (Brinkmann *et al.*, 1970; Rabideau, 1978; Dalling *et al.*, 1981; Arslan *et al.*, 2005).

The fourteen atoms of the anthracene skeleton in (I) (Fig. 1) have a total puckering amplitude $Q = 0.167(2) \text{ \AA}$ (Cremer & Pople, 1975). The methylene chain connecting the two rings




Figure 1
ORTEP drawing of (I) with the atom-numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 35% probability level. Unlabelled atoms are related to labelled atoms by $(-x, -y, 1-z)$.

exhibits an *anti-anti-anti* conformation. The O1–N–C10–C11 and O2–N–C10–C11 torsion angles (Table 1) indicate that the O atoms of the nitro groups lie in the same plane as the methylene chain.

Examination of the packing diagram (Fig. 2) reveals that the molecular packing is mainly determined by intermolecular C–H \cdots O interactions (Table 2).

Experimental

The title compound was obtained as the decomposition product of (*E,E*)-1,4-bis[9,10-dihydro-9-nitro-10(trinitromethyl)-9-anthryl]butane, which was synthesized by irradiation of a solution containing 21 mg (0.049 mmol) of 1,4-bis(9-anthryl)butane, 325 mg (1.67 mmol) of TNM, 45 ml pentane, and 5 ml CCl₄ under the conditions described by Arslan *et al.*, (2005). (*E,E*)-1,4-bis[9,10-dihydro-9-nitro-10(trinitromethyl)-9-anthryl]butane was obtained as yellow needles after removal of the solvents under reduced pressure and washing with acetone to remove the unreacted starting compound. The remaining yellow solid was column chromatographed using alumina (80–200 mesh, activity III) as the carrier and dichloromethane/hexane (1:4 *v/v*) as the eluent to yield the title compound (16.1% yield, m.p. 464–465 K, dichloromethane). Pale-yellow single crystals suitable for the X-ray diffraction study were grown from a concentrated solution of (I) in dichloromethane through slow evaporation under ambient conditions. ¹H-NMR (300 MHz, CDCl₃, p.p.m.): δ 8.33 [4 H, *d*, 2 \times (H1, H8)], 7.63 [8 H, *m*, 2 \times (H2, H3, H6, H7)], 7.34 [4 H, *d*, 2 \times (H4, H5)], 2.55 [4 H, *t*, 2 \times (H11a, H11b)], 0.38 [4 H, *m*, 2 \times (H12a, H12b)]

Crystal data

C₃₂H₂₄N₂O₆
 $M_r = 532.53$
 Monoclinic, $P2_1/c$
 $a = 11.316$ (1) Å
 $b = 8.330$ (1) Å
 $c = 13.880$ (2) Å
 $\beta = 97.728$ (9)°
 $V = 1296.6$ (2) Å³

$Z = 2$
 $D_x = 1.364$ Mg m⁻³
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 295$ (2) K
 Cube, pale yellow
 0.34 \times 0.34 \times 0.34 mm




Figure 2
The crystal packing of (I), viewed down the *b* axis. Dashed lines indicate C–H \cdots O interactions.

Data collection

Enraf-Nonius CAD-4
 diffractometer
 ω scans
 Absorption correction: none
 2296 measured reflections
 2296 independent reflections

1638 reflections with $I > 2\sigma(I)$
 $\theta_{\max} = 25.0^\circ$
 3 standard reflections
 frequency: 120 min
 intensity decay: 0.5%

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.110$
 $S = 1.01$
 2296 reflections
 181 parameters
 H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0433P)^2 + 0.4238P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.006$
 $\Delta\rho_{\max} = 0.19$ e Å⁻³
 $\Delta\rho_{\min} = -0.14$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

O1–N	1.199 (2)	O9–C9	1.224 (2)
O2–N	1.207 (2)	N–C10	1.562 (2)
O1–N–O2	123.27 (17)	C4A–C10–C10A	115.12 (15)
O1–N–C10	119.15 (16)	C4A–C10–N	105.39 (14)
O2–N–C10	117.58 (15)	C10A–C10–N	105.02 (14)
O1–N–C10–C11	1.4 (2)	O2–N–C10–C11	–178.02 (18)

Table 2

Hydrogen-bond geometry (Å, °).

<i>D</i> –H \cdots <i>A</i>	<i>D</i> –H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> –H \cdots <i>A</i>
C1–H1 \cdots O9 ⁱ	0.93	2.64	3.118 (3)	112

Symmetry code: (i) $-x + 1, -y, -z + 1$.

All H atoms were positioned geometrically and allowed to ride on their parent atoms at distances of 0.93 and 0.97 Å for aromatic and methylene H atoms, respectively, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Data collection: *CAD-4-PC* Software (Enraf-Nonius, 1993); cell refinement: *CAD-4-PC* Software; data reduction: *DATRD2* in *NRCVAX* (Gabe *et al.*, 1989); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1993); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors thank the Turkish Ministry of Education and the CSU College of Graduate Studies for their support of this work.

References

- Altomare, A., Casciarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343–350.
- Arslan, M., Baker, R. J., Masnovi, J. & Asker, E. (2005). *Acta Cryst.* **E61**, o4133–o4135.
- Brinkmann, A. W., Gordon, M., Harvey, R. G., Rabideau, P. W., Stothers, J. B. & Terney, A. L. (1970). *J. Am. Chem. Soc.* **92**, 5912–5916.
- Butts, C. P., Ebersson, L., Hartshorn, M. P., Robinson, W. T., Timmerman-Vaughan, D. J. & Young, D. A. W. (1996). *Acta Chem. Scand.* **50**, 29–47.
- Cox, A. (1998). *Photochemistry*, **29**, 164–203.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Dalling, D. K., Zilm, K. W., Grant, D. M., Heeschen, W. A., Horton, W. J. & Pugmire, R. J. (1981). *J. Am. Chem. Soc.* **103**, 4817–4824.
- Enraf-Nonius (1993). *CAD-4-PC Software*. Version 1.2. Enraf-Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
- Kochi, J. K. (1991). *Pure Appl. Chem.* **63**, 255–264.
- Lehnig, M. & Schürmann, K. (1998). *Eur. J. Org. Chem.* pp. 913–918.
- Rabideau, P. W. (1978). *Acc. Chem. Res.* **11**, 145–147.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.