

9-5-2013

# Adaptation of Scenedesmus Dimorphus to Brackish Water

Dustin Bowden  
*Cleveland State University*

Joanne M. Belovich  
*Cleveland State University, j.belovich@csuohio.edu*

## How does access to this work benefit you? Let us know!

Follow this and additional works at: [http://engagedscholarship.csuohio.edu/u\\_poster\\_2013](http://engagedscholarship.csuohio.edu/u_poster_2013)

 Part of the [Biomaterials Commons](#), and the [Plant Sciences Commons](#)

---

## Recommended Citation

Bowden, Dustin and Belovich, Joanne M., "Adaptation of Scenedesmus Dimorphus to Brackish Water" (2013). *Undergraduate Research Posters 2013*. Book 12.

[http://engagedscholarship.csuohio.edu/u\\_poster\\_2013/12](http://engagedscholarship.csuohio.edu/u_poster_2013/12)

This Article is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2013 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact [library.es@csuohio.edu](mailto:library.es@csuohio.edu).



This digital edition was prepared by MSL Academic Endeavors, the imprint of the Michael Schwartz Library at Cleveland State University.

# **Adaptation of *Scenedesmus dimorphus* to Brackish Water**

Fenn College of Engineering

Department of Chemical and Biomedical Engineering

**Student Researcher:** Dustin Bowden

**Faculty Advisor:** Joanne M. Belovich, Ph.D.

## **Abstract**

Microalgae is a promising biofuel feedstock for replacement of conventional transportation fuels. Microalgae does not require arable land for cultivation, and the biofuel production rate per acre of land is an order of magnitude greater than that needed for crop-based production methods. Though microalgae to biofuel processes are attractive, none have proven commercially successful due to the high costs of algae dewatering. Moreover, the scarcity of fresh water in many parts of the world prevents development of this process because of competition with drinking water supplies. Our lab has developed an efficient dewatering method using an inclined gravity settler. It is possible to adapt the freshwater *Scenedesmus dimorphus* to brackish water, without significantly changing growth rate nor lipid content. In this research, we investigate whether this saltwater-adapted algae species can be dewatered using the gravity settling method.

The settling velocity was measured by a settling column using absorbance at 600 nm to measure cell concentration. Our preliminary results yielded settling velocities of 0.76 cm/h, which is similar to earlier measurements of 0.87 cm/h for freshwater *S. dimorphus*. Based on these settling velocity measurements, we have found that saltwater acclimation will have minimal effect on the separation efficiency of *S. dimorphus*.