

9-4-2014

Alternative Reaction Pathways to Metformin Hydrochloride

Mohammed S. Suleiman
Cleveland State University

Stephen A. Reeves
Cleveland State University

Jorge E. Gatica
Cleveland State University, j.gatica@csuohio.edu

How does access to this work benefit you? Let us know!

Follow this and additional works at: http://engagedscholarship.csuohio.edu/u_poster_2014

 Part of the [Medicinal and Pharmaceutical Chemistry Commons](#)

Recommended Citation

Suleiman, Mohammed S.; Reeves, Stephen A.; and Gatica, Jorge E., "Alternative Reaction Pathways to Metformin Hydrochloride" (2014). *Undergraduate Research Posters 2014*. Book 36.
http://engagedscholarship.csuohio.edu/u_poster_2014/36

This Article is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2014 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.



This digital edition was prepared by MSL Academic Endeavors, the imprint of the Michael Schwartz Library at Cleveland State University.

Alternative Reaction Pathways to Metformin Hydrochloride

Washkewicz College of Engineering

Student Researchers: Mohammed S. Suleiman and Stephen A. Reeves

Faculty Advisor: Jorge E. Gatica

Abstract

Metformin Hydrochloride is an important pharmaceutical used for the treatment of type 2 diabetes. The current manufacturing of this product involves a well-known and proven process. The process includes the dissolution and reaction, followed by the precipitation of Metformin Hydrochloride. Although reliable and effective, the current process relies on the use of a solvent; which later needs to be eliminated from the precipitates. The purpose of this project is the investigation of an alternative reaction pathway which will avoid the use of solvents and simplify the final purification stage. The anticipated benefits include reduced costs for the processing and a final product which is closer to meet FDA and quality standards. These steps will eventually result in reducing the final market value of this important pharmaceutical. The investigation of different pathways was conducted using standard thermal characterization and surface analysis instrumentation. Namely, a differential scanning calorimeter (DSC) and a scanning electron microscope (SEM). The DSC was used for thermal characterization of the reactants and product standards; while the SEM was used to examine crystal morphology and elemental composition of reacting mixtures. Preliminary experiments were conducted using micro and laboratory scale solvent-less reacting environments. These experiments allowed identifying the presence of a single chemical reaction. The characterization results suggest that the alternative pathway can successfully synthesize Metformin Hydrochloride. Further characterization and testing protocols are currently being formulated.