5-15-2006

10,10'-Dinitro-10,10'-(propane-1,3-diyl)di-10H-anthracen-9-one

Mustafa Arslan
Sakarya University, Adapazari, Turkey

Erol Asker
Balıkesir University, Balıkesir, Turkey

John Masnovi
Cleveland State University, j.masnovi@csuohio.edu

Ronald J. Baker

Follow this and additional works at: https://engagedscholarship.csuohio.edu/scichem_facpub

Part of the [Organic Chemistry Commons](#)

[How does access to this work benefit you? Let us know!](#)

Recommended Citation

Arslan, M., Asker, E., Masnovi, J., & Baker, R. J. (2006). 10,10'-dinitro-10,10'-(propane-1,3-diyl)di-10H-anthracen-9-one. *Acta Crystallographica Section E*, 62(5), o2037-o2039. doi:10.1107/S1600536806013705

This Article is brought to you for free and open access by the Chemistry Department at EngagedScholarship@CSU. It has been accepted for inclusion in Chemistry Faculty Publications by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.

10,10'-Dinitro-10,10'-(propane-1,3-diyl)di-10H-anthracen-9-one

Mustafa Arslan,^{a*} Erol Asker,^b
John Masnovi^c and Ronald J.
Baker^c

^aSakarya Üniversitesi, Fen-Edebiyat Fakültesi
Kimya Bölümü, 54140 Esentepe/Adapazarı,
Turkey, ^bBalikesir Üniversitesi, Necatibey Eğitim
Fakültesi, 10100 Balikesir, Turkey, and
^cDepartment of Chemistry, Cleveland State
University, Cleveland, OH 44115, USA

Correspondence e-mail: marslan@sakarya.edu.tr

Key indicators

Single-crystal X-ray study

$T = 295\text{ K}$

Mean $\sigma(\text{C}-\text{C}) = 0.008\text{ Å}$

R factor = 0.076

wR factor = 0.214

Data-to-parameter ratio = 12.9

For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.

The title compound, $\text{C}_{31}\text{H}_{22}\text{N}_2\text{O}_6$, was obtained as the decomposition product of (*E,E*)-1,3-bis[9,10-dihydro-9-nitro-10-(trinitromethyl)-9-anthryl]propane, which was synthesized *via* a photochemical reaction of 1,3-di-9-anthrylpropane with tetranitromethane. Intermolecular $\text{C}-\text{H}\cdots\text{O}$ interactions are the most prominent features of the crystal packing; no indications of any intermolecular $\pi-\pi$ stacking were found.

Received 10 March 2006

Accepted 16 April 2006

Comment

Photonitration of various aromatic compounds using tetranitromethane (TNM) has attracted some attention as an alternative to the conventional nitration processes, which require the use of concentrated nitric and sulfuric acids (Kochi, 1991; Butts *et al.*, 1996; Cox, 1998; Lehnig & Schürmann, 1998). In general, addition of TNM to 9-alkyl-substituted anthracenes proceeds at the 9- and 10-positions, with the nitro group bonding to atom C9, bearing the alkyl group, and the trinitromethyl group being attached to the sterically less hindered unsubstituted C10 center. We have already reported the structure of the product of such a process, (*E*)-9,10-dihydro-9-methyl-9-nitro-10-(trinitromethyl)anthracene (Arslan *et al.*, 2005). One interesting feature of these trinitromethyl-substituted anthracene derivatives is that they contain the highly labile $\text{C}-\text{C}(\text{NO}_2)_3$ bond and, therefore, when passed through silica gel or an alumina column, easily decompose to form the corresponding anthrone derivatives. The decomposition process is believed to involve retro-aldol reaction.

Figure 1
ORTEP (Farrugia, 1997) drawing of (I), with the atom-numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 20% probability level. The H atoms are shown as a small circles of arbitrary radii.

Figure 2
The crystal packing of (I), viewed along the diagonal of the *bc* plane. Dashed lines indicate the C—H...O interactions.

In this paper we report the crystal structure of the title compound, (I), the decomposition product of (*E,E*)-1,3-bis[9,10-dihydro-9-nitro-10-(trinitromethyl)-9-anthryl]propane. The bond lengths and angles in the two anthracene ring systems (Table 1) are in agreement with each other, as well as with those of related compounds (Brinkmann *et al.*, 1970; Rabideau, 1978; Dalling *et al.*, 1981; Arslan *et al.*, 2005).

The 14 atoms of each anthracene system in (I) (Fig. 1) are coplanar to within 0.062 and 0.035 Å for the unprimed and primed ring systems, respectively. The trimethylene chain exhibits an *anti-anti* conformation. The dihedral angle formed by the anthracene planes is 22.51 (9)°.

Examination of the packing diagram (Fig. 2) reveals that the crystal packing is mainly determined by intermolecular C—H...O interactions; there are no indications of intermolecular π - π stacking in the crystal structure of (I).

Experimental

The title compound was obtained as the decomposition product of (*E,E*)-1,3-bis[9,10-dihydro-9-nitro-10-(trinitromethyl)-9-anthryl]-

propane, which was synthesized by irradiation with visual light of a solution containing 20 mg (0.050 mmol) of 1,3-bis(9-anthryl)propane, 325 mg (1.67 mmol) of TNM, 45 ml of pentane, and 5 ml of CCl₄ as described by Arslan *et al.* (2005). A 450 W Hanovia medium-pressure mercury lamp with a 500 nm sharp cut-off filter was used as a light source. The (*E,E*)-1,3-bis[9,10-dihydro-9-nitro-10-(trinitromethyl)-9-anthryl]propane was column chromatographed using alumina (80–200 mesh, activity III) as the carrier and dichloromethane–hexane as eluant to give (I) (38.9% yield, m.p. 464–465 K). Single crystals suitable for X-ray diffraction study were grown from a concentrated solution of (I) in dichloromethane through slow evaporation under ambient conditions.

Crystal data

C₃₁H₂₂N₂O₆
 $M_r = 518.51$
 Monoclinic, $P2_1/n$
 $a = 13.438$ (2) Å
 $b = 14.490$ (3) Å
 $c = 13.974$ (3) Å
 $\beta = 109.505$ (5)°
 $V = 2564.8$ (9) Å³

$Z = 4$
 $D_x = 1.343$ Mg m⁻³
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 295$ (2) K
 Block, yellow
 $0.32 \times 0.29 \times 0.13$ mm

Data collection

Enraf–Nonius CAD-4
 diffractometer
 ω scans
 Absorption correction: none
 4554 measured reflections
 4554 independent reflections

1912 reflections with $I > 2\sigma(I)$
 $\theta_{\max} = 25.1^\circ$
 3 standard reflections
 every 120 min
 intensity decay: 1.1%

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.076$
 $wR(F^2) = 0.214$
 $S = 0.99$
 4554 reflections
 352 parameters

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0908P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.20$ e Å⁻³
 $\Delta\rho_{\min} = -0.20$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

O1—N	1.187 (5)	O2'—N'	1.210 (5)
O2—N	1.175 (5)	O9'—C9'	1.221 (5)
O9—C9	1.228 (5)	N—C10	1.554 (5)
O1'—N'	1.185 (4)	N'—C10'	1.558 (5)
O1—N—O2	122.4 (4)	C4A—C10—N	105.5 (3)
O1—N—C10	117.6 (4)	C10A—C10—N	105.0 (3)
O2—N—C10	120.0 (4)	C11—C10—N	107.6 (3)
O1'—N'—O2'	123.4 (4)	C11'—C10'—C10A'	111.6 (4)
O1'—N'—C10'	118.7 (4)	C11'—C10'—C4A'	111.9 (3)
O2'—N'—C10'	117.6 (4)	C10A'—C10'—C4A'	115.5 (3)
C4A—C10—C10A	115.1 (3)	C11'—C10'—N'	107.9 (3)
C4A—C10—C11	111.2 (4)	C10A'—C10'—N'	105.2 (3)
C10A—C10—C11	111.7 (4)	C4A'—C10'—N'	104.0 (3)
O2—N—C10—C10A	−116.4 (6)	O1'—N'—C10'—C11'	−164.3 (4)
O1—N—C10—C11	−175.4 (4)	O2'—N'—C10'—C11'	21.6 (5)
N—C10—C11—C12	178.1 (4)	C11—C12—C11'—C10'	178.4 (4)
C10—C11—C12—C11'	179.1 (4)	N'—C10'—C11'—C12	178.5 (4)

Table 2
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C2'-H2'\cdots O2^i$	0.93	2.67	3.586 (10)	168
$C4-H4\cdots O9^{ii}$	0.93	2.60	3.417 (6)	147
$C5-H5\cdots O2^{iii}$	0.93	2.69	3.363 (7)	130
$C5'-H5'\cdots O2^{iii}$	0.93	2.53	3.343 (7)	146
$C7'-H7'\cdots O2^{iv}$	0.93	2.57	3.473 (7)	165
Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$; (ii) $x - \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$; (iii) $-x + 1, -y + 1, -z + 3$; (iv) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$				

All H atoms were positioned geometrically and allowed to ride on their corresponding parent atoms at C—H distances of 0.93 and 0.97 Å for aromatic and methylene H atoms, respectively, with $U_{iso}(H) = 1.2U_{eq}(\text{parent atom})$.

Data collection: *CAD-4-PC Software* (Enraf–Nonius, 1993); cell refinement: *CAD-4-PC Software*; data reduction: *DATRD2* in *NRCVAX* (Gabe *et al.*, 1989); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1993); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors thank the Turkish Ministry of Education and the CSU College of Graduate Studies for their support of this work.

References

- Altomare, A., Cascarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343–350.
- Arslan, M., Baker, R. J., Masnovi, J. & Asker, E. (2005). *Acta Cryst.* **E61**, o4133–o4135.
- Brinkmann, A. W., Gordon, M., Harvey, R. G., Rabideau, P. W., Stothers, J. B. & Terney, A. L. (1970). *J. Am. Chem. Soc.* **92**, 5912–5916.
- Butts, C. P., Eberson, L., Hartshorn, M. P., Robinson, W. T., Timmerman-Vaughan, D. J. & Young, D. A. W. (1996). *Acta Chem. Scand.* **50**, 29–47.
- Cox, A. (1998). *Photochemistry*, **29**, 164–203.
- Dalling, D. K., Zilm, K. W., Grant, D. M., Heeschen, W. A., Horton, W. J. & Pugmire, R. J. (1981). *J. Am. Chem. Soc.* **103**, 4817–4824.
- Enraf–Nonius (1993). *CAD-4-PC Software*. Version 1.2. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
- Kochi, J. K. (1991). *Pure Appl. Chem.* **63**, 255–264.
- Lehnig, M. & Schürmann, K. (1998). *Eur. J. Org. Chem.* pp. 913–918.
- Rabideau, P. W. (1978). *Acc. Chem. Res.* **11**, 145–147.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.