

Cleveland State University

EngagedScholarship@CSU

---

Undergraduate Research Posters 2015

Undergraduate Research Posters


---

2015

## Optimizing the design and potency of diterpenic acid derivatives to improve cell membrane permeability and Hsp27 targeting characteristics

Aicha Quamine  
*Cleveland State University*

Follow this and additional works at: [https://engagedscholarship.csuohio.edu/u\\_poster\\_2015](https://engagedscholarship.csuohio.edu/u_poster_2015)

 Part of the [Life Sciences Commons](#), [Medicine and Health Sciences Commons](#), and the [Physical Sciences and Mathematics Commons](#)

[How does access to this work benefit you? Let us know!](#)

---

### Recommended Citation

Quamine, Aicha, "Optimizing the design and potency of diterpenic acid derivatives to improve cell membrane permeability and Hsp27 targeting characteristics" (2015). *Undergraduate Research Posters 2015*. 19.

[https://engagedscholarship.csuohio.edu/u\\_poster\\_2015/19](https://engagedscholarship.csuohio.edu/u_poster_2015/19)

This Book is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2015 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact [library.es@csuohio.edu](mailto:library.es@csuohio.edu).



***Optimizing the design and potency of diterpenic acid derivatives to improve cell membrane permeability and Hsp27 targeting characteristics***

College of Sciences and Health Professions

**Student Researcher:** Aicha Quamine

**Faculty Advisor:** Bin Su

**Abstract**

Heat shock protein 27 (Hsp27) acts as a protective protein allowing it to play an integral role in aiding cancer cell resistance. Stress-induced Hsp 27 overexpression aids in the stabilization of partially denatured proteins to establish protein refolding resulting in thermotolerance, inhibition of apoptosis, cytoprotection, etc. As a result of rapid proliferation and general instability, cancer cells exhibit increased dependency on the support of Hsp 27 and its chaperone proteins, therefore generating an ideal target for anticancer therapy. Copalic acid, a clerodane diterpenoid, has already been confirmed in effective chaperone inhibition and antiproliferative synergistic effect. We hypothesize that by optimizing the structure of Copalic acid derivatives to increase solubility and drug potency we will increase anti-chaperone activity thereby hindering phosphorylation of Hsp27 affecting cell proliferation and stability. combinatorial chemistry strategies were used to develop and modify Copalic acid derivatives while synthesizing various dimers to enhance potency. Through utilizing NMR to identify synthesis alterations and MTT colorimetric assay we will continue to monitor and optimize the derived compounds.

*\*Supported by the McNair Scholars Program*