

Cleveland State University

EngagedScholarship@CSU

Undergraduate Research Posters 2012

Undergraduate Research Posters

9-6-2012

Obtaining a Pure Protein Using an ELP-Tagged TEV Protease

Teisha L. Mullins

Cleveland State University, T.L.MULLINS@csuohio.edu

Follow this and additional works at: https://engagedscholarship.csuohio.edu/u_poster_2012

 Part of the [Amino Acids, Peptides, and Proteins Commons](#)

[How does access to this work benefit you? Let us know!](#)

Recommended Citation

Mullins, Teisha L., "Obtaining a Pure Protein Using an ELP-Tagged TEV Protease" (2012). *Undergraduate Research Posters 2012*. 27.

https://engagedscholarship.csuohio.edu/u_poster_2012/27

This Book is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2012 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.



Obtaining a Pure Protein Using an ELP-Tagged TEV Protease

College of Sciences and Health Professions

Student Researcher: Teisha Mullins

Faculty Advisors: Kiril A. Streletzky, Ph.D., Department of Physics;
Nolan B Holland, Ph.D., Department of Chemical and Biochemical
Engineering

Abstract

Elastin-like polypeptides (ELPs) reversibly aggregate and phase separate above and solubilize below a specific transition temperature (T_t). ELPs are composed of repeats of the structural sequence of the elastin protein, Gly-Xaa-Gly-Val-Pro (GXGVP)_n. Xaa is a “guest residue” in the sequence and can be any amino acid, with the exception of proline, P, in order to alter the temperature response. ELPs can then be attached to recombinant target proteins as a tag to facilitate protein purification.

We present here use of GLGVP for tagging and purifying a gadolinium binding protein domain. Leucine is used as the guest residue in order to lower the T_t below room temperature, enabling purification to take place over a more manageable temperature range. After protein purification, it is ideal to remove the ELP tag from the target protein. To accomplish this, a tobacco etch virus (TEV) protease cleavage site was inserted between the target protein and the ELP tag. The TEV protease is also tagged with an ELP so that after proteolytic cleavage the ELP tag from the target protein, a single round of thermal cycling can remove both the free ELP tag and the ELP-tagged TEV protease, leaving the purified target protein.