

9-6-2012

Light Scattering Study of Elongated Particles: From Inorganic Nanorice to Polypeptide Micelles

Philip Dee

Cleveland State University, P.DEE@csuohio.edu

Follow this and additional works at: https://engagedscholarship.csuohio.edu/u_poster_2012

 Part of the [Laboratory and Basic Science Research Commons](#), and the [Optics Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

Dee, Philip, "Light Scattering Study of Elongated Particles: From Inorganic Nanorice to Polypeptide Micelles" (2012). *Undergraduate Research Posters 2012*. 40.

https://engagedscholarship.csuohio.edu/u_poster_2012/40

This Book is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2012 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.



This digital edition was prepared by MSL Academic Endeavors, the imprint of the Michael Schwartz Library at Cleveland State University.

Light Scattering Study of Elongated Particles: From Inorganic Nanorice to Polypeptide Micelles

College of Sciences and Health Professions

Student Researcher: Phil Dee

Faculty Advisors: Kiril A. Strelitzky, Ph.D., Department of Physics;
Nolan B Holland, Ph.D., Department of Chemical and Biochemical Engineering

Abstract

Utilizing the powerful experimental technique of Dynamic Light Scattering (DLS) for size characterization of anisotropic particles can be extremely misleading. Unfortunately, this point is often not realized by researchers who strive for particle sizing of nanoparticles in suspensions. The first goal of this study was to highlight the ambiguities of the DLS experiment on elongated particles. The second goal was to demonstrate the power of Depolarized Dynamic Light Scattering (DDLS) in probing the anisotropy of different types of nanoparticles. Both goals were realized by studying two very different systems: inorganic FeOOH nanorice and elastin like polypeptide (ELP) micelles. The difference between the two systems is fundamental as FeOOH particles are solid, contain no water, and, therefore, are easily imaged using SEM, TEM, and AFM. Polypeptide micelles are soft particles filled with water, and, therefore, not easily imaged by abovementioned techniques. Perfecting DDLS on a system like FeOOH would allow less ambiguous interpretation of light scattering experiments on ELP micelles. We present a consistent analysis of DDLS results on FeOOH nanorice and outline the potential difficulties and challenges of DDLS application for polypeptide micelles.