

Cleveland State University

EngagedScholarship@CSU

---

Undergraduate Research Posters 2016

Undergraduate Research Posters

---

2016

## Effect of Salt Concentration on the Composition of Elastin-Like Polypeptides in the Condensed Coacervate Phase

Louise L. Beckstrom  
*Cleveland State University*

Adam Maraschy  
*Cleveland State University*

Daniel P. Miller  
*Cleveland State University*

Follow this and additional works at: [https://engagedscholarship.csuohio.edu/u\\_poster\\_2016](https://engagedscholarship.csuohio.edu/u_poster_2016)

 Part of the [Engineering Commons](#)

[How does access to this work benefit you? Let us know!](#)

---

### Recommended Citation

Beckstrom, Louise L.; Maraschy, Adam; and Miller, Daniel P., "Effect of Salt Concentration on the Composition of Elastin-Like Polypeptides in the Condensed Coacervate Phase" (2016). *Undergraduate Research Posters 2016*. 47.

[https://engagedscholarship.csuohio.edu/u\\_poster\\_2016/47](https://engagedscholarship.csuohio.edu/u_poster_2016/47)

This Book is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2016 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact [library.es@csuohio.edu](mailto:library.es@csuohio.edu).



## ***Effect of Salt Concentration on the Composition of Elastin-Like Polypeptides in the Condensed Coacervate Phase***

Washkewicz College of Engineering

**Student Researchers:** Louise L. Beckstrom, Adam Maraschy, and Daniel P. Miller

**Faculty Advisor:** Nolan B. Holland

### **Abstract**

Elastin-Like Polypeptides (ELPs) are thermo-responsive polymers which could potentially be used as vehicles for drug delivery. The nanoparticle vehicles are called micelles and the basic structure is made of ELP-foldon. This ELP-foldon has a head group (foldon) that is hydrophilic and three tails (ELP) that are hydrophobic. Above a certain temperature, the transition temperature ( $T_t$ ), the hydrophobic tails aggregate together to form spheres with the hydrophilic headgroups on the outside. Inside the micelles, linear ELP and drugs can be captured. Although micelles can form without linear ELP, they are bigger when it is present. At lower temperatures the ELPs are fully soluble and exist as one solution phase, but at higher temperatures there exists two liquid phases. Even though they are both liquids, one is more viscous than the other. The less viscous phase is mostly water and the more viscous one is a condensed phase called a coacervate. Coacervate consists of a high concentration of protein, but still contains a significant amount of water. It appears as glue-like substance with a slight yellow tinge. The concentration of linear ELP may depend on the concentration of salt in solution and since the body naturally contains salt, knowing this relationship would be useful in designing micelles for drug delivery. We varied the concentration of salt in linear ELP to see its effect on the volume of the condensed coacervate at different temperatures.