Cleveland State University

EngagedScholarship@CSU

Undergraduate Research Posters 2018

Undergraduate Research Posters

2018

Effects of greater palatine nerve anesthesia on the activity of hyoid musculature during swallowing

Besa Bunjaku Cleveland State University

Follow this and additional works at: https://engagedscholarship.csuohio.edu/u_poster_2018



Part of the Medicine and Health Sciences Commons

How does access to this work benefit you? Let us know!

Recommended Citation

Bunjaku, Besa, "Effects of greater palatine nerve anesthesia on the activity of hyoid musculature during swallowing" (2018). Undergraduate Research Posters 2018. 52.

https://engagedscholarship.csuohio.edu/u_poster_2018/52

This Book is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2018 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.



Effects of greater palatine nerve anesthesia on the activity of hyoid musculature during swallowing

College of Sciences and Health Professions

Student Researcher: Besa Bunjaku

Faculty Advisor: Andrew Lammers

Abstract

Feeding is an important activity for all animals. An expansive array of sensory provide information to the brain about food handling during feeding. The brain then coordinates muscles to push the bolus into the esophagus, bypassing the airway. Previous work shows considerable coordination among sensor arrays and the nerves supplying them. Therefore it is likely that anesthetizing part of the oral cavity (in this case, the tissue covering the hard palate) will cause changes in the timing of muscles that are active during swallowing, even though these muscles are supplied by completely different nerves. We examined the timing of the mylohyoid (floor of the mouth) and thyrohyoid (a muscle that contracts during swallowing) via electromyography in three infant pigs while they drank milk from a bottle. Control data, with all sensory systems intact, were collected first, and then the tissue covering the hard palate was anesthetized. We used the electrical activity of the muscles to determine the timing of their contractions. We are still analyzing our data, but we expect to find that the muscles contract for longer periods of time, reflecting a more forceful push on the milk in the absence of sensory information to provide negative feedback.