

Cleveland State University

EngagedScholarship@CSU

---

Undergraduate Research Posters 2018

Undergraduate Research Posters


---

2018

## The role of synthesis solvent in particle size of metal organic frameworks

Jonathan Krisak  
*Cleveland State University*

Follow this and additional works at: [https://engagedscholarship.csuohio.edu/u\\_poster\\_2018](https://engagedscholarship.csuohio.edu/u_poster_2018)

 Part of the [Biomedical Engineering and Bioengineering Commons](#), and the [Chemical Engineering Commons](#)

[How does access to this work benefit you? Let us know!](#)

---

### Recommended Citation

Krisak, Jonathan, "The role of synthesis solvent in particle size of metal organic frameworks" (2018). *Undergraduate Research Posters 2018*. 63.

[https://engagedscholarship.csuohio.edu/u\\_poster\\_2018/63](https://engagedscholarship.csuohio.edu/u_poster_2018/63)

This Book is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2018 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact [library.es@csuohio.edu](mailto:library.es@csuohio.edu).



## ***Investigating Particle Size Manipulation for the Metal Organic Framework Cu-BTCs***

Washkewicz College of Engineering

**Student Researcher:** Jonathan Krisak

**Faculty Advisor:** Sasidhar Gumma

### **Abstract**

Metal organic frameworks are a class of nanoporous materials with pore sizes ranging from 0.5 to 3 nm and high surface areas (500-6000 m<sup>2</sup>/g). These materials have potential applications in industrial catalysis, separation and purification, bio-mimetics, drug delivery, semiconductors, sensors and other electronics. The aim of this study is to understand the role of solvent in control of the particle size of the final MOF product. CuBTC MOF has been used as a model MOF in this study to understand this effect.

Altering the dielectric constant of solvents is a potential method of controlling the particle size. The data obtained in this work depicts a direct correlation between the particle size and the dielectric constant of the solvent mixture. Deviations from this rule can be potentially explained by slow evaporation rate, longer nucleation growth, as found in literature, or instability of the hydroxide ions.

Literature states that the donor number and vapor pressure of the solvents also seem to affect the particle size. We observed that, while there is a direct correlation between particle size and donor number, no clear trend was observed between vapor pressure and particle size in this study.