

11-1-2016

Eventual Quasi-Linearity of The Minkowski Length

Ivan Soprunov

Cleveland State University, i.soprunov@csuohio.edu

Jenya Soprunova

Kent State University

Follow this and additional works at: https://engagedscholarship.csuohio.edu/scimath_facpub

 Part of the [Mathematics Commons](#)

[How does access to this work benefit you? Let us know!](#)

Repository Citation

Soprunov, Ivan and Soprunova, Jenya, "Eventual Quasi-Linearity of The Minkowski Length" (2016).

Mathematics Faculty Publications. 189.

https://engagedscholarship.csuohio.edu/scimath_facpub/189

This Article is brought to you for free and open access by the Mathematics and Statistics Department at EngagedScholarship@CSU. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.

Eventual quasi-linearity of the Minkowski length

Ivan Soprunov, Jenya Soprunova

0. Introduction

Let P be a d -dimensional lattice polytope in \mathbb{R}^d . Recall that the lattice diameter $\ell(P)$ is defined as one less than the largest number of collinear lattice points in P . The Minkowski length is a natural extension of this notion. For any $1 \leq n \leq d$, let $L_n(P)$ be the largest number of lattice polytopes of positive dimension whose Minkowski sum is at most n -dimensional and is contained in P . We call $L_n(P)$ the n th Minkowski length of P , and $L(P) = L_d(P)$ simply the Minkowski length of P . Note $L_1(P)$ coincides with the lattice diameter $\ell(P)$, as in this case the Minkowski summands are collinear

lattice segments. It is not hard to show (see the discussion after Definition 1.1) that $L_n(P)$ is the largest number of lattice segments whose Minkowski sum is at most n -dimensional and is contained in P .

The Minkowski length $L(P)$ of a lattice polytope $P \subset \mathbb{R}^d$ was first introduced in [7] in relation to studying parameters of toric surface codes. Every lattice polytope P defines a space $\mathcal{L}(P)$ of Laurent polynomials (over some field) whose monomials have exponent vectors lying in P . Such spaces naturally appear in the theory of toric varieties. The algebraic interpretation of the Minkowski length is the following: $L(P)$ is the largest number of irreducible factors a polynomial $f \in \mathcal{L}(P)$ may have. This information is particularly important when one studies zeros of polynomials in $\mathcal{L}(P)$, see [6,5,7,8,4]. A number of results concerning $L(P)$ appeared in [3,7,9].

Let $tP = \{tx \in \mathbb{R}^d \mid x \in P\}$ be the dilate of P by a positive integer factor t . The main result of this paper explains the behavior of $L_n(tP)$ as a function of the scaling factor $t \in \mathbb{N}$ in the spirit of the Ehrhart theory. In Theorem 2.20 we prove that for any lattice polytope P the function $L_n(tP)$ is eventually quasi-polynomial with linear constituents (we say “quasi-linear” for short), which contributes positively to the “ubiquitousness of quasi-polynomials” phenomenon declared by Kevin Woods [10]. For an introduction to the Ehrhart theory we refer the reader to the wonderful book by M. Beck and S. Robins [2].

To prove eventual quasi-linearity of the Minkowski lengths $L_n(P)$ we define and study their rational counterparts: a sequence of rational numbers $\lambda_1(P) \leq \dots \leq \lambda_d(P) = \lambda(P)$ associated with P . Here $\lambda_1(P)$ is the rational diameter of P and $\lambda_n(P)$ is the “asymptotic” Minkowski length, i.e. $\lambda_n(P) = \lim_{t \rightarrow \infty} L_n(tP)/t$. In Theorem 2.15 we prove that $\lambda_n(P) = L_n(kP)/k$ for some $k \in \mathbb{N}$.

Although an algorithm for computing $L(P)$ was presented in dimensions two and three (see [3,7]), there have been no explicit formulas for $L(P)$ even for simplices. Here we prove that $L(t\Delta) = t$ for any unimodular simplex Δ and any $t \in \mathbb{N}$. This result allowed us to write explicit answers for $L(P)$ for other classes of polytopes such as coordinate boxes and polytopes of degree one (see Corollary 2.2 and examples afterwards). In Section 3 we prove that for lattice triangles the Minkowski length coincides with the lattice diameter. The final part of the paper contains some examples and open questions.

1. Preliminaries

We start with some standard terminology from geometric combinatorics. A polytope $P \subset \mathbb{R}^d$ is called *lattice* (resp. *rational*) if its vertices have integer (resp. rational) coordinates. A vector $v \in \mathbb{Z}^d$ is called *primitive* if the greatest common divisor of its coordinates is 1. A lattice segment is called *primitive* if it contains exactly two lattice points. A d -dimensional simplex is called *unimodular* if its vertices affinely generate the lattice \mathbb{Z}^d .

Given a lattice polytope $P \subset \mathbb{R}^d$, denote by $\text{Vol}_d(P)$ the Euclidean d -dimensional volume of P . Note that the d -dimensional volume of any parallelepiped formed by a basis of \mathbb{Z}^d equals 1. More generally, suppose P is contained in an n -dimensional rational affine subspace $a+H$ for a rational linear subspace $H \subset \mathbb{R}^d$ and $a \in \mathbb{Q}^d$. Denote by $\text{Vol}_n(P)$ the n -dimensional volume of P normalized such that the n -dimensional volume of any parallelepiped formed by a basis of the lattice $H \cap \mathbb{Z}^d$ equals 1.

1.1. Minkowski length

Let P and Q be convex polytopes in \mathbb{R}^d . Their *Minkowski sum* is the set

$$P + Q = \{p + q \in \mathbb{R}^d \mid p \in P, q \in Q\},$$

which is again a convex polytope.

Definition 1.1. Let P be a lattice polytope in \mathbb{R}^d . Define the *Minkowski length* $L = L(P)$ of P to be the largest number of lattice polytopes Q_1, \dots, Q_L of positive dimension whose Minkowski sum is contained in P . Any such sum $Q_1 + \dots + Q_L$ is called a *maximal decomposition* in P .

We refer the reader to [3] for examples illustrating this definition. It is clear from the definition that $L(P)$ is monotone with respect to inclusion: $L(P) \leq L(Q)$ if $P \subseteq Q$, and is superadditive with

respect to the Minkowski sum: $L(P + Q) \geq L(P) + L(Q)$. Also, $L(P)$ is invariant under unimodular transformations (isomorphisms of the lattice \mathbb{Z}^d).

There is a natural partial order on the set of maximal decompositions in P , as defined in [3]. Namely, we say that

$$Q_1 + \dots + Q_L \leq P_1 + \dots + P_L$$

if $Q_1 + \dots + Q_L$ is contained in $P_1 + \dots + P_L$ after a possible lattice translation. Minimal elements with respect to this partial order are called *smallest maximal decompositions*. Clearly, every smallest maximal decomposition is the Minkowski sum of L lattice segments, i.e. is a lattice zonotope.

Any lattice (resp. rational) zonotope Z can be written in the form

$$Z = a + \alpha_1[0, v_1] + \dots + \alpha_m[0, v_m], \tag{1.1}$$

for some $m \in \mathbb{N}$, distinct primitive vectors $v_i \in \mathbb{Z}^d$, positive integer (resp. rational) numbers α_i , and a lattice (resp. rational) point $a \in \mathbb{R}^d$. In this case, we set

$$|Z| := \alpha_1 + \dots + \alpha_m.$$

The following result from [3] gives a universal bound for the number of distinct summands in a smallest maximal decomposition.

Proposition 1.2 ([3]). *Let $P \subset \mathbb{R}^d$ be a lattice polytope. Then every smallest maximal decomposition in P has at most $2^d - 1$ distinct summands.*

1.2. Quasi-polynomials

Here we recall the definition of a quasi-polynomial function.

Definition 1.3. A function $f : \mathbb{N} \rightarrow \mathbb{Q}$ is called a *quasi-polynomial* if there exist $k \in \mathbb{N}$ and polynomials $p_0, \dots, p_{k-1} \in \mathbb{Q}[t]$, called the *constituents* of f , such that

$$f(t) = p_r(t) \quad \text{whenever } t \equiv r \pmod k, \text{ for } 0 \leq r \leq k.$$

The smallest such k is called the *period* of f . If all the constituents of f are linear we say that f is *quasi-linear*. Finally, we say $f : \mathbb{N} \rightarrow \mathbb{Q}$ is *eventually quasi-linear* if $f(t)$ coincides with a quasi-linear function for all large enough t .

Example 1.4. The function $f(t) = 3 \lfloor \frac{t}{3} \rfloor + 4$ is quasi-linear with period $k = 3$, where $\lfloor x \rfloor$ denotes the floor of x . Indeed,

$$f(t) = \begin{cases} t + 4, & \text{if } t \equiv 0 \pmod 3 \\ t + 3, & \text{if } t \equiv 1 \pmod 3 \\ t + 2, & \text{if } t \equiv 2 \pmod 3. \end{cases}$$

2. Main theorems

Before we prove our main result about eventual quasi-linearity of the Minkowski length, we will look at some instances when it is, in fact, linear. The simplest such example is when $P = \Delta$, a unimodular d -simplex.

Theorem 2.1. *Let Δ be a unimodular d -simplex and $t \in \mathbb{N}$. Then*

$$L(t\Delta) = tL(\Delta) = t.$$

Proof. After a unimodular transformation we may assume that Δ is the standard d -simplex, i.e. the convex hull of $\{0, e_1, \dots, e_d\}$ where e_i are the standard basis vectors. First, it is easy to see that $L(\Delta) = 1$. Also, $L(t\Delta) \geq t$ as $t\Delta$ contains the Minkowski sum of t lattice segments $[0, e_1]$.

We prove the converse by induction on d . The case $d = 1$ is trivial. Denote $L = L(t\Delta)$. Let Z be a smallest maximal decomposition in $t\Delta$ and let $a \in Z$ be a vertex with the smallest sum of the coordinates, which we denote by α . We have

$$Z = a + [0, v_1] + \cdots + [0, v_L],$$

where $v_i \in \mathbb{Z}^d$ are primitive, not necessarily distinct vectors. Note that the sum of the coordinates of each v_i is non-negative, by the choice of a . Suppose the first k of the vectors v_1, \dots, v_L have the sum of the coordinates equals zero, for $0 \leq k \leq L$. Then the subzonotope

$$Z' = a + [0, v_1] + \cdots + [0, v_k]$$

is contained in $\alpha\Delta'$, where Δ' is the facet of Δ whose points have the sum of the coordinates equal to 1. This implies $k \leq L(\alpha\Delta')$. By induction $L(\alpha\Delta') = \alpha$, hence, $k \leq \alpha$. Now the point $v = a + v_{k+1} + \cdots + v_L$ lies in Z and has the sum of the coordinates at least $\alpha + L - k \geq L$. On the other hand, v lies in $t\Delta$, so its sum of the coordinates is at most t . Therefore, $L \leq t$. \square

Corollary 2.2. *Let P be a lattice polytope contained in $\alpha\Delta$ for some unimodular simplex Δ and $\alpha \in \mathbb{N}$. If P contains the Minkowski sum of α lattice segments then $L(P) = \alpha$. Consequently, $L(tP) = tL(P)$.*

Example 2.3. Let Π be a lattice coordinate box in \mathbb{R}^d , i.e. $\Pi = [0, \alpha_1 e_1] \times \cdots \times [0, \alpha_d e_d]$, where e_i are the standard basis vectors and α_i are non-negative integers. Clearly, $\Pi = \alpha_1[0, e_1] + \cdots + \alpha_d[0, e_d]$ and Π is contained in $(\alpha_1 + \cdots + \alpha_d)\Delta_d$, where Δ_d is the standard d -simplex. Therefore,

$$L(\Pi) = \alpha_1 + \cdots + \alpha_d.$$

We also have $L(t\Pi) = tL(\Pi)$.

Example 2.4. According to a result of Batyrev and Nill [1], a d -dimensional polytope P has *degree one* if and only if P is either

- (1) the $d - 2$ iterated pyramid over the triangle $\Delta_2 = \text{Conv}\{(0, 0), (2, 0), (0, 2)\}$, or
- (2) the $d - n$ iterated pyramid over a Lawrence prism Q defined by a sequence of integers $0 < h_1 \leq \cdots \leq h_n$:

$$Q = \text{Conv}\{0, e_1, \dots, e_{n-1}, e_1 + h_1 e_n, \dots, e_{n-1} + h_{n-1} e_n, h_n e_n\} \subset \mathbb{R}^n. \quad (2.1)$$

Corollary 2.2 implies that in the first case $L(P) = 2$ since P contains the segment $[0, 2e_1]$ and $P \subset 2\Delta_d$. In the second case

$$L(P) = \begin{cases} h_n, & \text{if } h_{n-1} < h_n \\ h_n + 1, & \text{if } h_{n-1} = h_n. \end{cases}$$

Indeed, if $h_{n-1} < h_n$ then $P \subset h_n \Delta_d$ and P contains the segment $[0, h_n e_n]$. If $h_{n-1} = h_n$ then $P \subset (h_n + 1)\Delta_d$ and P contains the rectangle $[0, e_{n-1}] + [0, h_n e_n]$.

2.1. Rational Minkowski length

Let P be an arbitrary lattice polytope in \mathbb{R}^d . The following is a generalization of Definition 1.1.

Definition 2.5. Let P be a lattice polytope in \mathbb{R}^d . Define the n th Minkowski length $L = L_n(P)$ of P to be the largest number of lattice polytopes Q_1, \dots, Q_L of positive dimension whose Minkowski sum is at most n -dimensional and is contained in P .

Clearly, $L_1(P) \leq \cdots \leq L_{d-1}(P) \leq L_d(P) = L(P)$. Note $L_1(P)$ coincides with the *lattice diameter* $\ell(P)$, which is defined as one less than the largest number of collinear lattice points in P .

Example 2.6. Let \square be the unit square in \mathbb{R}^2 . Then

$$L_1(\square) = 1 \quad \text{and} \quad L_2(\square) = L(\square) = 2.$$

For any unimodular d -simplex Δ and any $t \in \mathbb{N}$ we have $L_1(t\Delta) \geq t$ as $t\Delta$ contains the segment $[0, te_1]$. By Theorem 2.1, $L(t\Delta) = t$, hence

$$L_1(t\Delta) = \cdots = L_d(t\Delta) = L(t\Delta) = t.$$

If P is the $d - n$ iterated pyramid over a Lawrence prism Q as in (2.1) then $L_1(P) = h_n$ and $L_2(P) = \cdots = L_d(P) = L(P)$.

The following is a rational analog of the Minkowski length.

Definition 2.7. The number

$$\lambda(P) = \sup_{t \in \mathbb{N}} \frac{L(tP)}{t}$$

is called the *rational Minkowski length* of P . More generally, the n th rational Minkowski length of P is

$$\lambda_n(P) = \sup_{t \in \mathbb{N}} \frac{L_n(tP)}{t}.$$

The following proposition asserts that the numbers $\lambda_n(P)$ are well-defined.

Proposition 2.8. For any $t \in \mathbb{N}$ we have $L_n(tP) \leq t\alpha$, where $\alpha \in \mathbb{N}$ is such that $P \subseteq \alpha\Delta$ for a unimodular simplex Δ .

Proof. It is enough to consider the case $n = d$. Then it is immediate from Theorem 2.1: $L(tP) \leq L(t\alpha\Delta) = t\alpha$. \square

It follows from the definition that $\lambda_1(P) \leq \cdots \leq \lambda_{d-1}(P) \leq \lambda_d(P) = \lambda(P)$.

Remark 2.9. As $L_n(tP)$ satisfies the superadditivity property, the supremum in the above definition may be replaced with the limit, by Fekete's lemma. We will not be using this result in our further discussion.

Definition 2.10. Let $K \subset \mathbb{R}^d$ be a rational polytope. For any primitive vector $v \in \mathbb{Z}^d$ define $s_v(K)$ to be the largest rational number s such that the segment $[0, sv]$ is contained in K after a translation by a rational vector. The *rational diameter* $s(K)$ is the maximum of $s_v(K)$ over all primitive $v \in \mathbb{Z}^d$.

It is not hard to see that $s(P) = \lambda_1(P)$ for any lattice polytope P . Indeed, for any $t \in \mathbb{N}$, the polytope P contains a segment $a + [0, sv]$ for some $a \in \mathbb{Q}^d$, primitive $v \in \mathbb{Z}^d$, and $s = L_1(tP)/t$. Thus, $\lambda_1(P) \leq s(P)$. Conversely, if $a + [0, s(P)v]$ is contained in P for some $a \in \mathbb{Q}^d$ and primitive $v \in \mathbb{Z}^d$ then there exists $t \in \mathbb{N}$ such that $ta + [0, ts(P)v]$ is a lattice segment contained in tP , i.e. $s(P) \leq L_1(tP)/t \leq \lambda_1(P)$. As a corollary, we obtain $L_1(P) = \lfloor \lambda_1(P) \rfloor$, as $\ell(P) = \lfloor s(P) \rfloor$.

In our main theorem below (Theorem 2.15) we show that $\lambda(P)$ as well as all $\lambda_n(P)$ are, in fact, rational numbers. First, we need a few lemmas.

Lemma 2.11. Let K be a convex body in \mathbb{R}^d and fix $\varepsilon > 0$. Then the set

$$U_\varepsilon(K) = \{v \in \mathbb{Z}^d \mid v \text{ primitive, } s_v(K) \geq \varepsilon\}$$

is finite.

Proof. First, note that if $K \subseteq K'$ then $s_v(K) \leq s_v(K')$, and $s_v(\alpha K) = \alpha s_v(K)$ for $\alpha \in \mathbb{Q}$. Thus it is enough to prove the statement for $K = \mathbb{B}$, the d -dimensional unit ball. Let $v \in \mathbb{Z}^d$ be primitive. By definition $s_v(\mathbb{B})$ is the number $s \in \mathbb{Q}$ such that $\|sv\| = 2$, where $\|\cdot\|$ is the usual Euclidean norm. It follows that $s_v(\mathbb{B}) \geq \varepsilon$ if and only if $\|v\| \leq 2/\varepsilon$. In other words, $U_\varepsilon(\mathbb{B})$ is a lattice set contained in the ball of radius $2/\varepsilon$ and so is finite. \square

Lemma 2.12. Let $Z = a + \alpha_1[0, v_1] + \cdots + \alpha_m[0, v_m]$ be a smallest maximal decomposition in P and $n = \dim Z$. Then for any $1 \leq i_1 < \cdots < i_n \leq m$ the n -dimensional volume of the parallelepiped formed by v_{i_1}, \dots, v_{i_n} is no greater than n^d .

Proof. We may assume that v_{i_1}, \dots, v_{i_n} are linearly independent. Let $a + H$ be the affine span of Z and let $\mathbb{L} = H \cap \mathbb{Z}^d$ be the corresponding lattice of rank n . It is well-known that the n -dimensional volume of the parallelepiped formed by n linearly independent lattice vectors $w_1, \dots, w_n \in \mathbb{L}$ equals the number of lattice points in the half-open parallelepiped

$$\{\lambda_1 w_1 + \cdots + \lambda_n w_n \mid 0 \leq \lambda_i < 1 \text{ for } 1 \leq i \leq n\},$$

which is less than the number of lattice points in the closed parallelepiped.

Let $\{v_{i_1}, \dots, v_{i_n}\}$ be a subset of the set of vectors appearing in the decomposition Z . We claim that the parallelepiped Π they form has at most n^d lattice points. Indeed, consider the image of $\Pi \cap \mathbb{Z}^d$ in $(\mathbb{Z}/n\mathbb{Z})^d$. If there are two lattice points in Π congruent mod $(n\mathbb{Z})^d$ then the lattice segment E containing them lies in Π and has lattice length n . Therefore, if we replace $[0, v_{i_1}] + \cdots + [0, v_{i_n}]$ in the decomposition Z with E we obtain a maximal decomposition Z' properly contained in Z . This contradicts the fact that Z is a smallest maximal decomposition. This shows that all lattice points of Π are different in $(\mathbb{Z}/n\mathbb{Z})^d$, i.e. their number cannot exceed n^d . \square

Lemma 2.13. Let $B = \{u_1, \dots, u_n\}$ be a basis for a rational linear subspace $H \subseteq \mathbb{R}^d$, and fix a constant N . Let $\text{Vol}_n(u_1, \dots, v, \dots, u_n)$ denote the n -dimensional volume of the parallelepiped formed by u_1, \dots, u_n where u_i is replaced with a vector v . Then the set

$$V(B) = \{v \in H \cap \mathbb{Z}^d \mid \text{Vol}_n(u_1, \dots, v, \dots, u_n) \leq N, \text{ for all } 1 \leq i \leq n\}$$

is finite.

Proof. Fix a coordinate system in H by choosing a basis for the lattice $H \cap \mathbb{Z}^d$. Write $v = x_1 u_1 + \cdots + x_n u_n$ for some $x_i \in \mathbb{R}$. Then by Cramer's rule

$$|x_i| = \frac{\text{Vol}_n(u_1, \dots, v, \dots, u_n)}{\text{Vol}_n(u_1, \dots, u_n)} \leq \frac{N}{\text{Vol}_n(u_1, \dots, u_n)} =: c_i.$$

Therefore, the set $V(B)$ is contained in the set of lattice points of the parallelepiped formed by $\{\pm c_1 u_1, \dots, \pm c_n u_n\}$, and, hence, is finite. \square

Lemma 2.14. Let $P \subset \mathbb{R}^d$ be a lattice polytope. Fix an ordered collection of primitive vectors $\mathbf{v} = (v_1, \dots, v_m) \in (\mathbb{Z}^d)^m$. Then the set of zonotopes

$$\mathcal{Z}(\mathbf{v}) = \{Z = a + \alpha_1[0, v_1] + \cdots + \alpha_m[0, v_m] \mid a \in \mathbb{R}^d, \alpha_i \in \mathbb{R}_{\geq 0}, Z \subseteq P\}$$

is a rational polytope in \mathbb{R}^{d+m} . The function $|\cdot| : \mathcal{Z}(\mathbf{v}) \rightarrow \mathbb{R}, Z \mapsto |Z|$ is an integer linear function on $\mathcal{Z}(\mathbf{v})$.

Proof. With every such zonotope Z we associate a point $z = (a, \alpha_1, \dots, \alpha_m) \in \mathbb{R}^{d+m}$. Note that Z is the convex hull of the following set of 2^m points in \mathbb{R}^d :

$$K = \left\{ a + \sum_{i \in I} \alpha_i v_i \mid I \subseteq \{1, \dots, m\} \right\}.$$

Clearly $Z \subseteq P$ if and only if $K \subset P$ which is expressed by 2^m rational linear inequalities in $d + m$ variables. Therefore, they define a rational polytope $\mathcal{Z}(\mathbf{v})$ in \mathbb{R}^{d+m} (the boundedness of $\mathcal{Z}(\mathbf{v})$ follows from that of P).

The function $|\cdot| : \mathcal{Z}(\mathbf{v}) \rightarrow \mathbb{R}, Z \mapsto |Z|$ is determined by the sum of the last m coordinates in \mathbb{R}^{d+m} , hence, is an integer linear function on $\mathcal{Z}(\mathbf{v})$. \square

Notice that reordering of the v_i does not change the zonotope Z , so the polytope $\mathcal{Z}(\mathbf{v})$, as well as the function $|\cdot| : \mathcal{Z}(\mathbf{v}) \rightarrow \mathbb{R}$, is invariant under permutations of the last m coordinates.

Now we are ready for our main result.

Theorem 2.15. *Let $P \subset \mathbb{R}^d$ be a lattice polytope. Then*

$$\lambda(P) = \frac{L(kP)}{k},$$

for some $k \in \mathbb{N}$.

Proof. Consider the polytope tP for some $t \in \mathbb{N}$. It follows from Proposition 1.2, that tP contains a smallest maximal decomposition Z with $m \leq M := 2^d - 1$ distinct summands

$$Z = a + \alpha_1[0, v_1] + \cdots + \alpha_m[0, v_m], \quad (2.2)$$

where $a \in \mathbb{Z}^d$, $v_i \in \mathbb{Z}^d$ are primitive, and α_i are positive integers whose sum equals the Minkowski length $L(tP)$. Therefore, P contains a rational zonotope

$$Z/t = a/t + (\alpha_1/t)[0, v_1] + \cdots + (\alpha_m/t)[0, v_m]$$

with $|Z/t| = L(tP)/t$. Conversely, every rational zonotope Z in P has the form

$$Z = a + \alpha_1[0, v_1] + \cdots + \alpha_m[0, v_m], \quad (2.3)$$

for some $a \in \mathbb{Q}^d$, primitive $v_i \in \mathbb{Z}^d$, and non-negative rationals α_i . Then there exists $t \in \mathbb{N}$ such that tZ is a lattice zonotope in tP , and so $|Z| \leq L(tP)/t \leq \lambda(P)$. Therefore, $\lambda(P)$ is the supremum of the function $Z \mapsto |Z|$ on the set of all rational zonotopes Z contained in P .

We will show below that there exist $\delta > 0$, independent of t , and a finite set of primitive vectors $V_\delta \subset \mathbb{Z}^d$ satisfying the following property: If Z is a smallest maximal decomposition in tP for some $t \in \mathbb{N}$ then

$$\lambda(P) - \delta < |Z/t| \leq \lambda(P)$$

implies that v_1, \dots, v_m lie in V_δ . By Lemma 2.14, $\lambda(P)$ equals the maximum of the linear function $Z \mapsto |Z|$ on the union of rational polytopes $\mathcal{Z}(\mathbf{v})$ over all collections $\mathbf{v} = (v_1, \dots, v_m) \in (V_\delta)^m$, hence, $\lambda(P) = |Z'|$ for some rational zonotope $Z' \subset P$. Choose $k \in \mathbb{N}$ such that kZ' is a lattice zonotope in kP . Then

$$\lambda(P) = \frac{L(kP)}{k},$$

as required.

It remains to prove the existence of $\delta > 0$ and V_δ satisfying the above property. Denote $\lambda = \lambda(P)$, and $\lambda_n = \lambda_n(P)$, the n th rational Minkowski length of P . Let $e \geq 1$ be the smallest integer such that $\lambda_n = \lambda(P)$ for all $n \geq e$. Then to find λ it is enough to consider only smallest maximal decompositions in tP of dimension at most e . Let Z be such a decomposition, as in (2.2).

The case $e = 1$ is easy—we set $\delta = \lambda/2$ and $V_\delta = U_\delta(P)$, as in Lemma 2.11, which is a finite set. In this case $|Z/t| = \alpha_1/t \leq s_{v_1}(P) < \delta = \lambda/2$, unless $v_1 \in V_\delta$.

If $e > 1$ we have

$$\lambda_1 \leq \cdots \leq \lambda_{e-1} < \lambda_e = \cdots = \lambda_d = \lambda.$$

Set $\delta = (\lambda - \lambda_{e-1})/2$ and choose

$$0 < \varepsilon < \min \left\{ \frac{\lambda - \delta}{M}, \frac{\delta}{M - e} \right\}.$$

If no v_i lies in $U_\varepsilon(P)$ then $\alpha_i/t \leq s_{v_i}(P) < \varepsilon$, and so $|Z/t| \leq m\varepsilon < \lambda - \delta$. Thus, we may assume that $v_i \in U_\varepsilon(P)$ for $1 \leq i \leq k$ and $v_i \notin U_\varepsilon(P)$ for $k < i \leq m$.

First, suppose that $\{v_1, \dots, v_k\}$ spans an e -dimensional subspace. By Lemmas 2.12 and 2.13, there are only finitely many choices for each v_i for $k < i \leq m$. Thus we define V_δ to be the union of $U_\varepsilon(P)$

and the finite sets $V(B)$ for every subset $B = \{u_1, \dots, u_e\} \subset U_\varepsilon(P)$ which spans an e -dimensional subspace.

Next, suppose the dimension of the span of $\{v_1, \dots, v_k\}$ is less than e . We may assume that v_l, \dots, v_m lie outside of this span for some $k < l \leq m$. Then we have

$$|Z/t| \leq \lambda_{e-1} + (\alpha_l/t) + \dots + (\alpha_m/t) < \lambda_{e-1} + (m-l)\varepsilon.$$

By the choice of ε , and since $l > e$, the latter is smaller than $\lambda - \delta$. \square

Remark 2.16. The same arguments as above show that for any $1 \leq n \leq d$,

$$\lambda_n(P) = \frac{L(k_n P)}{k_n},$$

for some $k_n \in \mathbb{N}$. In particular, all $\lambda_n(P)$ are rational numbers.

2.2. Quasi-linearity of the Minkowski length

The result of Theorem 2.15 allows us to make the following definition.

Definition 2.17. The smallest $k \in \mathbb{N}$ satisfying $\lambda(P) = L(kP)/k$ is called the *period* of P .

In Theorem 2.20 we prove that the Minkowski length is eventually quasi-linear, but first we are going to show that the rational Minkowski length is linear (Proposition 2.19). We will need the following lemma.

Lemma 2.18. Let k be the period of P . Then $L(tP) = t\lambda(P)$ whenever $k \mid t$, $t \in \mathbb{N}$.

Proof. Since tP contains the Minkowski sum of t/k copies of kP , we have

$$L(tP) \geq (t/k)L(kP) = t\lambda(P).$$

On the other hand, $t\lambda(P) \geq L(tP)$, by the definition of $\lambda(P)$. \square

Proposition 2.19. For any $t \in \mathbb{N}$ we have $\lambda(tP) = t\lambda(P)$.

Proof. We have

$$\lambda(tP) = \max_{s \in \mathbb{N}} \frac{L(stP)}{s} = t \max_{s \in \mathbb{N}} \frac{L(stP)}{st} \leq t\lambda(P).$$

On the other hand, by Theorem 2.15 and Lemma 2.18,

$$t\lambda(P) = \frac{kt\lambda(P)}{k} = \frac{L(ktP)}{k} \leq \lambda(tP),$$

where k is the period of P . \square

Theorem 2.20. Let P be a lattice polytope in \mathbb{R}^d with period k . Then the function $L(tP)$ is eventually quasi-linear with period at most k . More explicitly, there exist integers c_r , for $0 \leq r < k$, such that $L(rP) \leq c_r \leq r\lambda(P)$ and

$$L(tP) = k\lambda(P) \left\lfloor \frac{t}{k} \right\rfloor + c_r$$

whenever $t \equiv r \pmod{k}$ and t is large enough.

Proof. Fix $0 \leq r < k$ and let $t \equiv r \pmod k$. Denote $c_r(t) = L(tP) - k\lambda(P) \lfloor \frac{t}{k} \rfloor$. Note that $c_r(t)$ is an integer as $k\lambda(P) = L(kP)$. We will show that $c_r(t)$ is constant for t large enough. Indeed, $c_r(t)$ are bounded from above:

$$c_r(t) = L(tP) - k\lambda(P) \left\lfloor \frac{t}{k} \right\rfloor \leq t\lambda(P) - k\lambda(P) \left\lfloor \frac{t}{k} \right\rfloor = r\lambda(P).$$

Also they increase:

$$c_r(t+k) = L(tP+kP) - k\lambda(P) \left\lfloor \frac{t+k}{k} \right\rfloor \geq L(tP) + L(kP) - k\lambda(P) \left\lfloor \frac{t+k}{k} \right\rfloor = c_r(t),$$

where we used $L(kP) = k\lambda(P)$ in the last equality. Therefore, the integers $c_r(t)$ for $t \equiv r \pmod k$ eventually stabilize to a constant c_r .

We have already seen that $c_r \leq r\lambda(P)$. For the other inequality, let $t = qk + r$. Then, using Lemma 2.18, we obtain

$$c_r(t) = L(tP) - qk\lambda(P) \geq L(qkP) + L(rP) - qk\lambda(P) = L(rP). \quad \square$$

Remark 2.21. The above proof works just as well if we replace $L(P)$ with $L_n(P)$ for any $1 \leq n \leq d$ and apply Remark 2.16. Therefore, each n th Minkowski length of P is eventually quasi-linear with period at most k_n . Since $L_1(P) = \lfloor \lambda_1(P) \rfloor$, the function $L_1(tP)$ is, in fact, quasi-linear.

3. Dimension two

In this section we deal with lattice polytopes in dimension two. We prove an upper bound on the rational length of P in terms of other well-known invariants of P —the Euclidean area and the lattice width of P . As an application we give a formula for $\lambda(tP)$ and $L(tP)$ for any triangle P in \mathbb{R}^2 .

Let $P \subset \mathbb{R}^d$ be a lattice polytope and $v \in \mathbb{Z}^2$ a primitive vector. Recall that the *lattice width of P in the direction v* is the integer $\max_{x \in P} \langle x, v \rangle - \min_{x \in P} \langle x, v \rangle$. Here $\langle x, v \rangle$ is the standard inner product in \mathbb{R}^d . The smallest lattice width over all primitive $v \in \mathbb{Z}^d$ is called the *lattice width of P* and is denoted $w(P)$.

Proposition 3.1. *Let $P \subset \mathbb{R}^2$ be a lattice polygon. Then $\lambda(P) \leq \frac{2\text{Vol}_2(P)}{w(P)}$ where $\text{Vol}_2(P)$ is the Euclidean area and $w(P)$ the lattice width of P .*

Proof. By the proof of Theorem 2.15, $\lambda(P) = |Z|$ for some rational zonotope $Z \subseteq P$ with at most 3 distinct summands, i.e.

$$Z = a + \alpha_1[0, v_1] + \alpha_2[0, v_2] + \alpha_3[0, v_3],$$

where $v_i \in \mathbb{Z}^2$ are distinct primitive vectors, $a \in \mathbb{Q}^2$, and $\alpha_i \in \mathbb{Q}$. We have $|Z| = \alpha_1 + \alpha_2 + \alpha_3$.

Let w_i be the lattice width of P in the direction of a primitive vector v_i^\perp , orthogonal to v_i . We claim that

$$\alpha_1 w_1 + \alpha_2 w_2 + \alpha_3 w_3 \leq 2 \text{Vol}_2(P). \quad (3.1)$$

Indeed, let A_i (resp. B_i) be a vertex of P where the inner product $\langle x, v_i^\perp \rangle$ attains its minimum (resp. maximum). Similarly, let E_i (resp. I_i) be the side of Z where $\langle x, v_i^\perp \rangle$ attains its minimum (resp. maximum). Connect A_i to E_i and B_i to I_i for $i = 1, 2, 3$ by line segments. Also triangulate Z (if it is not one-dimensional) by drawing the diagonals through the center of Z . We obtain a (not necessarily convex) triangulated polygon S inside P , see Fig. 1.

Note that the sum of the areas of the four triangles with bases E_i and I_i equals $\frac{1}{2}\alpha_i w_i$. Therefore, the left hand side of (3.1) represents twice the area of S , and (3.1) follows.

It remains to note that $\alpha_1 w_1 + \alpha_2 w_2 + \alpha_3 w_3 \geq (\alpha_1 + \alpha_2 + \alpha_3)w(P) = \lambda(P)w(P)$. \square

Below we apply this bound to give an explicit formula for the (rational) Minkowski length of any triangle. Recall the *lattice diameter* $\ell(P)$ and the *rational diameter* $s(P)$ defined in Section 2.1.




Fig. 1. A triangulated polygon inside P .

Corollary 3.2. Let $T \subset \mathbb{R}^2$ be a lattice triangle. Let $s(T)$ be its rational diameter and $\ell(T)$ its lattice diameter. Then

$$\lambda(T) = s(T) \quad \text{and} \quad L(T) = \lfloor s(T) \rfloor = \ell(T).$$

Consequently, $\lambda(tT) = s(T)t$ and $L(tT) = \lfloor s(T)t \rfloor$.

Proof. Let $v \in \mathbb{Z}^2$ be a primitive vector such that the lattice width of T in the direction orthogonal to v equals $w(T)$. Then $s_v(T)w(T) = 2 \text{Vol}_2(T)$, where $s_v(T)$ as in Definition 2.10. It follows that $s_v(T)$ is, in fact, $s(T)$. Applying Proposition 3.1, we get $\lambda(T) \leq s(T)$. Conversely, T contains the segment E parallel to $[0, s(T)v]$. Therefore, by the proof of Theorem 2.15, $\lambda(T) \geq |E| = s(T)$.

As for $L(T)$, it is clear that $L(T) \leq \lambda(T) = s(T)$, by the definition of $\lambda(P)$. Thus, $L(T) \leq \lfloor s(T) \rfloor$. On the other hand, T contains a translation of the lattice segment $[0, \lfloor s(T) \rfloor v]$, hence $L(T) \geq \lfloor s(T) \rfloor$.

Finally, by above, $\lambda(tT) = s(tT) = ts(T)$ and $L(tT) = \lfloor s(tT) \rfloor = \lfloor ts(T) \rfloor$. \square

Remark 3.3. The above proof shows that our bound in Proposition 3.1 is tight, as $\lambda(T) = s(T) = 2 \text{Vol}_2(T)/w(T)$ for any lattice triangle T .

4. Examples and open problems

In this section we illustrate our results with several examples and raise some questions.

Our first example shows that $L(tP)$ can have an arbitrarily large period k .

Example 4.1. Let $T_k \subset \mathbb{R}^2$ denote the triangle with vertices $(0, 0)$, $(k, 1)$, and $(1, k)$, for $k \geq 2$. It is not hard to see that $\ell(T_k) = k - 1$ and $s(T_k) = k - 1/k$. By Corollary 3.2,

$$L(tT_k) = \left\lfloor \left(k - \frac{1}{k}\right)t \right\rfloor,$$

which is a quasi-linear function with period k .

Example 4.2. Let P be a square with vertices $(2, 0)$, $(3, 2)$, $(1, 3)$, and $(0, 1)$. One readily sees that $\text{Vol}_2(P) = 5$, $w(P) = 3$, and $L(P) = 3$. Therefore, by Proposition 3.1, $\lambda(P) \leq 10/3$.

Note that $3P$ contains a zonotope Z with $|Z| = 10$ (in fact, Z is a square with vertices $(2, 2)$, $(7, 2)$, $(2, 7)$, and $(7, 7)$). Therefore, $10 \leq L(3P) \leq 3\lambda(P) \leq 10$, which implies that $\lambda(P) = 10/3$ and $L(tP)$ has period $k = 3$. By Lemma 2.18, if $t = 3q$ then $L(tP) = 10q$. Now if $t = 3q + 1$ we have

$$10q + 3 = L(3qP) + L(P) \leq L(tP) \leq 10t/3 = 10q + 10/3,$$

and, hence, $L(tP) = 10q + 3$. Similarly, $L(tP) = 10q + 6$ when $t = 3q + 2$. Therefore,

$$L(tP) = 10 \left\lfloor \frac{t}{3} \right\rfloor + 3r, \quad \text{if } t \equiv r \pmod{3}.$$

Looking at the above examples one may suspect that $L(P) = \lfloor \lambda(P) \rfloor$ for any polytope P . This would imply that $L(tP)$ is not just eventually quasi-linear, but quasi-linear:

$$L(tP) = k\lambda(P) \left\lfloor \frac{t}{k} \right\rfloor + L(rP),$$

for any $t \equiv r \pmod{k}$ (see Theorem 2.20).

However, $L(P) = \lfloor \lambda(P) \rfloor$ does not hold even for the case of lattice polygons, as demonstrated by the following example.

Example 4.3. Let $P = 2Q$ where Q is the square with vertices $(1, 0)$, $(5, 1)$, $(4, 5)$, and $(0, 4)$. Then by Proposition 3.1,

$$\lambda(P) = 2\lambda(Q) \leq \frac{68}{5}.$$

Also $L(5P)$ contains Z with $|Z| = 68$ (namely, Z is the square with vertices $(8, 8)$, $(8, 42)$, $(42, 42)$, and $(42, 8)$), hence, $\lambda(P) = 68/5$. By observation, $L(P) = 12$, which illustrates that $\lambda(P) - L(P)$ can be as large as $8/5$.

Problem 1. Find the supremum of $\lambda(P) - L(P)$ over all lattice polytopes $P \subset \mathbb{R}^d$.

It is not hard to see that $\lambda(P) - L(P) < 4$ for any lattice polygon $P \subset \mathbb{R}^2$, but we are confident that this bound could be improved.

In all 2-dimensional examples we computed, the function $L(tP)$ was always quasi-linear. Although we do not expect this to be the case in general, we have not been able to produce a counterexample.

Problem 2. Prove that $L(tP)$ is quasi-linear or give an example of a lattice polytope P for which $L(tP)$ is not quasi-linear.

Finally, we have seen that $L(P) = L_1(P)$ when P is a positive integer dilate of a unimodular simplex as well as when P is any simplex in dimension two. This prompts the following problem.

Problem 3. Prove or disprove that for any simplex in \mathbb{R}^d its Minkowski length coincides with its lattice diameter.

Acknowledgment

The first author is partially supported by NSA Grant H98230-13-1-0279.

References

- [1] V. Batyrev, B. Nill, Multiples of lattice polytopes without interior lattice points, *Mosc. Math. J.* 7 (2007) 195–207. 349.
- [2] Matthias Beck, Sinai Robins, *Computing the Continuous Discretely: Integer-point Enumeration in Polyhedra*, in: Undergraduate Texts in Mathematics, Springer, 2007.
- [3] O. Beckwith, M. Grimm, J. Soprunova, B. Weaver, Minkowski length of 3D lattice polytopes, *Discrete Comput. Geom.* 48 (4) (2012) 1137–1158.
- [4] V. Gauthier Umana, M. Velasco, Dual toric codes and polytopes of degree one, *SIAM J. Discrete Math.* 29 (1) (2015) 683–692.
- [5] J. Little, Remarks on generalized toric codes, in: *Finite Fields and their Applications*, Vol. 24, 2013, pp. 1–14.
- [6] J. Little, H. Schenck, Toric surface codes and Minkowski sums, *SIAM J. Discrete Math.* 20 (4) (2006) 999–1014 (electronic).
- [7] I. Soprunov, J. Soprunova, Toric surface codes and Minkowski length of polygons, *SIAM J. Discrete Math.* 23 (1) (2009) 384–400.
- [8] I. Soprunov, J. Soprunova, Bringing toric codes to the next dimension, *SIAM J. Discrete Math.* 24 (2) (2010) 655–665.
- [9] Josh Whitney, A bound on the minimum distance of three dimensional toric codes (Ph.D. Thesis), 2010.
- [10] Kevin Woods, The unreasonable ubiquitousness of quasi-polynomials, *Electron. J. Combin.* 21 (2014) P1.44.