

2004

Residues and Tame Symbols on Toroidal Varieties

Ivan Soprunov

Cleveland State University, i.soprunov@csuohio.edu

Follow this and additional works at: https://engagedscholarship.csuohio.edu/scimath_facpub

 Part of the [Mathematics Commons](#)

[How does access to this work benefit you? Let us know!](#)

Repository Citation

Soprunov, Ivan, "Residues and Tame Symbols on Toroidal Varieties" (2004). *Mathematics Faculty Publications*. 274.

https://engagedscholarship.csuohio.edu/scimath_facpub/274

This Article is brought to you for free and open access by the Mathematics and Statistics Department at EngagedScholarship@CSU. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.

Residues and tame symbols on toroidal varieties

Ivan Soprounov

ABSTRACT

We introduce a new approach to the study of a system of algebraic equations in $(\mathbb{C}^\times)^n$ whose Newton polytopes have sufficiently general relative positions. Our method is based on the theory of Parshin’s residues and tame symbols on toroidal varieties. It provides a uniform algebraic explanation of the recent result of Khovanskii on the product of the roots of such systems and the Gel’fond–Khovanskii result on the sum of the values of a Laurent polynomial over the roots of such systems, and extends them to the case of an algebraically closed field of arbitrary characteristic.

1. Introduction

1.1 The classical residue formula says that the sum of the residues of a rational 1-form ω over all points of a complex projective curve X is zero:

$$\sum_{x \in X} \operatorname{res}_x \omega = 0.$$

The standard proof of this formula uses Stokes’ theorem. In ‘Algebraic groups and class fields’ J.-P. Serre gives a purely algebraic proof of the residue formula which works over any algebraically closed field even of positive characteristic [Ser88].

In class field theory the residue formula has a multiplicative cousin, Weil’s reciprocity law, which states that the product of the tame symbols of any two rational functions f_0 and f_1 over all points of a projective curve X is one [Ser88]:

$$\prod_{x \in X} \langle f_0, f_1 \rangle_x = 1.$$

In the 1970s A. Parshin constructed higher-dimensional class field theory where he generalized the residue and the tame symbol [Par75]. Given an n -dimensional algebraic variety X and a rational form ω on X , Parshin defines the residue $\operatorname{res}_F \omega$ at each complete flag $F : X_0 \subset X_1 \subset \cdots \subset X_{n-1} \subset X$ of irreducible subvarieties of X . Similarly, given any $n + 1$ rational functions f_0, \dots, f_n on X , he defines the tame symbol $\langle f_0, \dots, f_n \rangle_F$ at each such flag F . Parshin’s residue and symbol satisfy not one but many reciprocity laws: Fix all subvarieties in the flag F except one, say X_i . Then the sum of the residues (product of the symbols) over all possible irreducible subvarieties X_i that can appear in the i th slot of F is zero (one) (Theorem A.4 in the Appendix).

The aim of the present paper is to look at some recent results of the theory of Newton polytopes from the point of view of this general theory of Parshin. More specifically, consider a system of Laurent polynomial equations in the n -torus:

$$f_1(t) = \cdots = f_n(t) = 0, \quad t \in (\mathbb{C}^\times)^n. \tag{1}$$

Suppose the Newton polytopes $\Delta_1, \dots, \Delta_n$ of the f_i have sufficiently general relative positions (see Definition 5.1). Then the system has a finite number of roots. O. Gel'fond and A. Khovanskii proved the following result [GK96].

THEOREM A. *The sum of the values of a Laurent polynomial f_0 over the roots of (1) counting multiplicities is equal to*

$$\sum_A (-1)^n c(A) \operatorname{res}_A \left(f_0 \frac{df_1}{f_1} \wedge \dots \wedge \frac{df_n}{f_n} \right),$$

where the sum is over the vertices A of $\Delta = \Delta_1 + \dots + \Delta_n$, $\operatorname{res}_A(f_0 df_1/f_1 \wedge \dots \wedge df_n/f_n)$ is the residue at a vertex (an explicit rational function in the coefficients of the f_i), and $c(A)$ is the combinatorial coefficient (an integer that reflects the combinatorial structure of the polytope Δ near the vertex A).

This is, in fact, a particular case of their residue formula [GK02] for the sum of the Grothendieck residues of a rational n -form which is regular in $(\mathbb{C}^\times)^n \setminus \{f_1 \dots f_n = 0\}$, over the roots of the system (1). The proof of the residue formula is topological and uses toric compactifications.

The following generalized Vieta formula for the product of roots of (1) has been obtained by Khovanskii [Kho99].

THEOREM B. *The product of the values of a Laurent monomial f_0 over the roots of (1) counting multiplicities is equal to*

$$\prod_A [f_0, \dots, f_n]_A^{(-1)^n c(A)},$$

where the product is over the vertices A of $\Delta = \Delta_1 + \dots + \Delta_n$, $[f_0, \dots, f_n]_A$ is the symbol at a vertex (an explicit Laurent monomial in the coefficients of the f_i), and $c(A)$ is the combinatorial coefficient.

The proof of this theorem uses the polyhedral homotopy method and regular subdivisions of polytopes.

The relation between Theorems A and B appears to be the same as the one between the one-dimensional residue formula and Weil's reciprocity. Moreover, the number $[f_0, \dots, f_n]_A$ is defined similarly to Parshin's tame symbol. This gives a motivation to search for a uniform explanation of these results in terms of the theory of residues and tame symbols.

The main obstruction to this is the notion of combinatorial coefficients $c(A)$ since they are defined as the local degrees of certain real non-algebraic maps. In the present paper we give an explicit algebraic description for the combinatorial coefficients as a signed number of certain complete flags of faces of Δ , thus putting them in the framework of Parshin's theory. (A similar description of the combinatorial coefficient was obtained by O. Gel'fond [Gel96] for some special collections of polytopes.) We provide a uniform algebraic proof of Theorems A and B based on Parshin's theory for toroidal varieties. We also extend them to the case of an arbitrary algebraically closed field.

1.2 The material of the paper is organized as follows. In § 2 we give an explicit formula for the degree of a map of polyhedral sets defined by some combinatorial data. As an application we obtain a new formula for the combinatorial coefficient.

In §§ 3 and 4 we consider Parshin's theory for toroidal pairs. A toroidal pair (X, D) consists of a normal variety X and a codimension-1 subvariety D such that, locally at each point, X is analytically isomorphic to an affine toric variety X_σ where the branches of D correspond to the invariant divisors of X_σ . We define residue and tame symbol at each point $x \in X$ for which the corresponding affine variety X_σ has a zero-dimensional orbit. This generalizes the notions of residue and symbol at a vertex from Theorems A and B. Our definition is similar to the one

of Parshin, but does not involve any particular choice of a complete flag. Using the algebraic description of the combinatorial coefficients we prove general results about symbol and residue on toroidal pairs (Theorems 3.15 and 4.8). In § 5 we show how these results imply Theorems A and B for arbitrary algebraically closed fields.

Finally, in the Appendix we include the definition of Parshin's residue and tame symbol and formulate the higher-dimensional reciprocity laws.

1.3 Remarks

There is a topological construction for the tame symbol based on Deligne's proof of Weil's reciprocity (see [BM96]). This construction provides a uniform topological explanation of the product of roots formula and the residue formula. Our approach is algebraic and works over algebraically closed fields of arbitrary characteristic.

A different formula for the product of the roots of a system (1) can be derived from Poisson's formula for the mixed resultant due to Pedersen and Sturmfels [PS93]. In this formula the product of the values of a monomial $f_0 = ct^m$ over the roots is represented by the product of the mixed resultant of f_0, \dots, f_n (which in this case is just c to the power of the mixed volume of $\Delta_1, \dots, \Delta_n$) and the facet resultants to certain powers. The assumption on the Newton polytopes of (1) implies that each facet resultant is a monomial in one of the coefficients of the system. To give an idea of how this formula is related to the one in Theorem B, let us assume that $f_0 = c$, for $c \neq 0, 1$. Then Poisson's formula gives the inductive formula for the mixed volume (e.g. see [BZ88, p. 166]), whereas Theorem B gives Khovanskii's formula for the mixed volume in terms of combinatorial coefficients [Kho99].

Parshin's residue is closely related to the toric residue defined by D. Cox [Cox96]. For different applications of residues in toric geometry we refer the reader to works of E. Cattani, D. Cox, A. Dickenstein, and B. Sturmfels [CCD97, CD97, CDS98].

1.4 In this paper k is always an algebraically closed field. A variety is a reduced separated scheme of finite type over k ; a subvariety is a reduced subscheme. By \mathbb{T} we denote the algebraic n -torus over k , $\mathbb{T} = (k^\times)^n$, and $M = \text{Hom}_{\text{alg. gp}}(\mathbb{T}, k^\times)$ the abelian rank- n group of characters of \mathbb{T} . Finally, X_σ denotes the affine toric variety $\text{Spec } k[\sigma \cap M]$ defined by a convex rational polyhedral cone σ in $M \otimes \mathbb{R}$.

2. Degree of polyhedral maps and combinatorial coefficient

In this section we show how to compute the degree of a map between two polyhedral sets which is defined by a map of the partially ordered sets of their faces. As an application we obtain an explicit combinatorial formula for the combinatorial coefficient.

2.1 Polyhedral maps

A polyhedral set is a finite union of convex compact polytopes intersecting in faces. We will assume that all the polytopes are embedded in a Euclidean space E of some big dimension. Then a polyhedral set is a topological space with the topology inherited from E . The dimension of a polyhedral set is the maximum of dimensions of the polytopes it contains. A polyhedral set is oriented if every polytope it contains is oriented.

Let X be a polyhedral set and $\mathcal{F}(X)$ the set of all faces of all polytopes appearing in X . The set $\mathcal{F}(X)$ is a finite partially ordered set by inclusion.

Consider two polyhedral sets X and Y , and fix a map $\psi : \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$ that preserves the partial ordering. A continuous piecewise linear map $f_\psi : X \rightarrow Y$ is called a *polyhedral map associated with ψ* if $f_\psi(G) \subset \psi(G)$ for every face $G \in \mathcal{F}(X)$.

Given any map $\psi : \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$ there exists a polyhedral map $f_\psi : X \rightarrow Y$. Moreover all such maps are homotopy equivalent within the class of all polyhedral maps associated with ψ .

PROPOSITION 2.1. *Each map $\psi : \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$ that respects the partial ordering defines a homotopy class of polyhedral maps $f_\psi : X \rightarrow Y$ associated with ψ .*

Proof. First, for every map of partially ordered sets $\psi : \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$ we construct a continuous piecewise linear map $f_\psi : X \rightarrow Y$ as follows.

Fix barycentric subdivisions of X and Y . Note that for any polyhedral set X there is a one-to-one correspondence between the set of all simplices in a barycentric subdivision of X and the set of all chains in $\mathcal{F}(X)$. Consider a k -simplex Δ^k in the subdivision of X . It corresponds to a chain $X_0 \subset \cdots \subset X_k$ in $\mathcal{F}(X)$. Let $\psi(X_0) \subseteq \cdots \subseteq \psi(X_k)$ be its image. It corresponds to a unique simplex (possibly of smaller dimension) in the subdivision of Y , the simplex being denoted by $\psi(\Delta^k)$. Since there is a unique linear map between two simplices that maps vertices of one simplex to the prescribed vertices of the other simplex, we get a map $f_\psi : X \rightarrow Y$ that sends each simplex Δ^k to the corresponding simplex $\psi(\Delta^k)$. Clearly this map agrees on the common faces of simplices of the subdivision and, hence, is continuous piecewise linear. By construction, $f_\psi(G) \subset \psi(G)$ for any $G \in \mathcal{F}(X)$.

Now suppose f_ψ and f'_ψ are two polyhedral maps associated with ψ . Then for each $0 \leq t \leq 1$ the map $f_\psi^t = (1-t)f_\psi + tf'_\psi$ is also associated with ψ . Indeed, fix a face $G \in \mathcal{F}(X)$. Then every point $x \in G$ is mapped to a point $f_\psi^t(x)$ on the segment joining $f_\psi(x)$ and $f'_\psi(x)$. Since both $f_\psi(x)$ and $f'_\psi(x)$ belong to the face $\psi(G)$, then $f_\psi^t(x)$ also does. Therefore $f_\psi^t(G) \subset \psi(G)$. \square

2.2 Flags and degree of polyhedral maps

Consider an oriented polyhedral set X . Let $\mathcal{X} : X_0 \subset \cdots \subset X_n$, $\dim X_i = i$ be a complete flag in X , i.e. a maximal chain of elements of $\mathcal{F}(X)$. With the flag \mathcal{X} we associate an ordered set of vectors (e_1, \dots, e_n) , where e_i begins at X_0 and points strictly inside X_i . Define the *sign* of \mathcal{X} to be 1 if (e_1, \dots, e_n) gives a positive oriented frame for X_n and -1 otherwise. It is easy to see that the sign does not depend on the choice of vectors e_1, \dots, e_n . We denote it by $\text{sgn } \mathcal{X}$.

Now consider a map of partially ordered sets $\psi : \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$, where X and Y are n -dimensional oriented polyhedral sets. Every polyhedral map $f_\psi : X \rightarrow Y$ associated with ψ induces a map of the n th homology groups:

$$H_n(f_\psi) : H_n(X) \rightarrow H_n(Y).$$

By Proposition 2.1 this map is the same for all choices of f_ψ . We call it the *degree* map of ψ . We will be concerned with the case when both groups $H_n(X)$ and $H_n(Y)$ are isomorphic to \mathbb{Z} . (This is true, for example, when X and Y are the boundaries of $(n+1)$ -dimensional polytopes.) Then the degree map is the multiplication by an integer, which we denote by $\deg(\psi)$. In the next theorem we show how to compute $\deg(\psi)$ as a signed number of certain complete flags in X .

Let $\mathcal{X} : X_0 \subset \cdots \subset X_n$ and $\mathcal{Y} : Y_0 \subset \cdots \subset Y_n$ be complete flags in X and Y , respectively. We will write $\psi(\mathcal{X}) = \mathcal{Y}$ if and only if $\psi(X_i) = Y_i$ for all $0 \leq i \leq n$. Define the *preimage* of \mathcal{Y} under ψ to be the set of all \mathcal{X} such that $\psi(\mathcal{X}) = \mathcal{Y}$.

THEOREM 2.2. *Let X and Y be two polyhedral sets as above, and $\psi : \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$ a map of partially ordered sets of their faces. Fix any complete flag \mathcal{Y} in Y . Then the degree of ψ is equal to the sign of \mathcal{Y} times the signed number of all complete flags \mathcal{X} in X in the preimage of \mathcal{Y} under ψ :*

$$\deg(\psi) = \text{sgn } \mathcal{Y} \sum_{\psi(\mathcal{X})=\mathcal{Y}} \text{sgn } \mathcal{X}.$$

Proof. By Proposition 2.1 we can choose any function in the homotopy class defined by ψ . We take f_ψ to be the piecewise linear function constructed in the proof of Proposition 2.1 using barycentric subdivisions of X and Y . We view f_ψ as a simplicial map between two simplicial complexes.

Fix any positive oriented n -simplex Δ_Y^n in the barycentric subdivision of Y . Then the degree of f_ψ is the number of all n -simplices Δ_X^n in X that are mapped to Δ_Y^n ; each simplex being counted with either sign plus or sign minus according to its orientation. Recall that the n -simplex Δ_Y^n corresponds to a complete flag \mathcal{Y} in Y and every n -simplex Δ_X^n corresponds to some complete flag \mathcal{X} in X . Clearly, the orientation of Δ_X^n coincides with the sign of the corresponding flag \mathcal{X} , and $f_\psi(\Delta_X^n) = \Delta_Y^n$ if and only if $\psi(\mathcal{X}) = \mathcal{Y}$. It remains to notice that if we fix a negative oriented Δ_Y^n then the number we obtain is the negative degree of f_ψ . \square

2.3 Combinatorial coefficient

The combinatorial coefficient is a local analog of the degree considered above.

Let $\sigma \subset \mathbb{R}^n$ be a convex polyhedral n -dimensional cone with apex A . Consider an ordered collection $D = (D_1, \dots, D_m)$ of m distinct non-empty closed subsets of σ , where $m \leq n$ and each set D_i is a union of facets of σ . Assume that they cover the boundary of σ and, if $m = n$, that the apex A is the only face of σ which is covered by all of them:

$$\partial\sigma = D_1 \cup \dots \cup D_m, \quad \text{if } m = n \text{ then } D_1 \cap \dots \cap D_n = \{A\}. \quad (2)$$

A continuous map $g : \sigma \rightarrow \mathbb{R}^n$ is called a *characteristic map* of the covering (2) if for each i , $1 \leq i \leq n$, the i th component g_i of g is non-negative and vanishes precisely on those faces of σ that belong to D_i . It is easy to see that all characteristic maps send the boundary of σ to the boundary of the positive octant \mathbb{R}_+^n such that $g^{-1}(0) \subseteq \{A\}$, and they are homotopy equivalent within the class of such maps.

DEFINITION 2.3. The local degree of the germ at A of the restriction of a characteristic map to the boundary of σ ,

$$\bar{g} : (\partial\sigma, A) \rightarrow (\partial\mathbb{R}_+^n, 0),$$

is called the *combinatorial coefficient* of the covering (2).

Clearly, the combinatorial coefficient is zero unless $m = n$. In the case when $m = n$, Theorem 2.2 provides us with a description of the combinatorial coefficient as the number of certain complete flags of faces of σ , counted with signs.

For a cone $\sigma \subset \mathbb{R}^n$ we let $\mathcal{F}(\partial\sigma)$ denote the partially ordered by inclusion set of the proper faces of σ . With a covering (2) we associate a map $\phi : \mathcal{F}(\partial\sigma) \rightarrow \mathcal{F}(\partial\mathbb{R}_+^n)$ by putting

$$\phi(\tau) = \mathbb{R}_+^n \cap \{y_{i_1} = \dots = y_{i_k} = 0\}$$

if and only if τ is a common face of D_{i_1}, \dots, D_{i_k} for $1 \leq i_l \leq n$, and k is maximal. Here (y_1, \dots, y_n) is a coordinate system for \mathbb{R}^n .

For any complete flag $\gamma_0 \subset \dots \subset \gamma_{n-1}$ in $\partial\mathbb{R}_+^n$ define its *preimage* under ϕ as the set of all complete flags $\sigma_0 \subset \dots \subset \sigma_{n-1}$ in $\partial\sigma$ such that $\phi(\sigma_i) = \gamma_i$, $0 \leq i \leq n-1$. Note that the preimage of any flag under ϕ is empty if $m < n$.

Fix the standard orientation of \mathbb{R}^n . We orient the boundary of every n -dimensional cone in \mathbb{R}^n in accordance with this fixed orientation. As before, define the *sign* of a complete flag $\sigma_0 \subset \dots \subset \sigma_{n-1}$ to be 1 if it gives a positive oriented frame for σ_{n-1} , and -1 otherwise.

THEOREM 2.4. *The combinatorial coefficient of a covering (2) is equal to the signed number of all complete flags in the preimage of any positive complete flag under ϕ .*

In particular, if $m = n$ the combinatorial coefficient is equal to the signed number of all complete flags $\sigma_0 \subset \cdots \subset \sigma_{n-1} \subset \sigma$, where σ_i is a common face of D_{i+1}, \dots, D_n of dimension i .

Proof. The case $m < n$ is obvious, so we assume that $m = n$. To be able to apply Theorem 2.2 we ‘compactify’ the cones σ and \mathbb{R}_+^n . Consider a pyramid with the vertex A and base D_0 which is a cross-section of σ by a generic hyperplane. Let X be the oriented boundary of the pyramid. Next consider the standard n -dimensional simplex defined in \mathbb{R}^n as the convex hull of the origin and the endpoints of the standard basis vectors. Let Y be its oriented boundary.

The subsets D_1, \dots, D_n of σ along with the base D_0 form a covering of X . Define the map $\psi : \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$, by putting $\psi(G) = \{y_{i_1} = \cdots = y_{i_k} = 0\}$ if and only if G is a common face of D_{i_1}, \dots, D_{i_k} for $0 \leq i_l \leq n$, and k is maximal. Here (y_0, y_1, \dots, y_n) are the barycentric coordinates for the simplex.

Note that the restriction of a characteristic map $g : \sigma \rightarrow \mathbb{R}^n$ to X defines a polyhedral map $f_\psi : X \rightarrow Y$ associated with ψ . According to Theorem 2.2, the degree of ψ is equal to the number of complete flags of faces of X counted with signs in the preimage under ψ of any positive complete flag of faces of Y . For example, one can take the flag

$$\{y_1 = \cdots = y_n = 0\} \subset \{y_2 = \cdots = y_n = 0\} \subset \cdots \subset \{y_n = 0\}. \quad \square$$

Remark 2.5. Notice that since there are $n!$ complete flags in \mathbb{R}_+^n we obtain $n!$ formulae for the combinatorial coefficient. If $m = n$ a choice of a complete flag corresponds to an order of D_1, \dots, D_n , and thus we can say that the combinatorial coefficient is skew-symmetric in D_1, \dots, D_n .

3. Toroidal symbol

The toroidal symbol is a slight modification of Parshin’s tame symbol for toroidal varieties. More precisely, consider a pair (X, D) consisting of a normal variety X and a codimension-1 subset D such that, in a formal neighborhood of each point, X is isomorphic to an affine toric variety X_σ and D corresponds to the invariant divisor $X_\sigma \setminus \mathbb{T}$. We distinguish special points on X for which the corresponding toric variety has a zero-dimensional orbit. At each such point $x \in X$ the toroidal symbol associates a non-zero element $[f_0, \dots, f_n]_x$ of the base field to every collection of $n + 1$ rational functions f_0, \dots, f_n on X with divisors in D .

Suppose the irreducible components of D are divided into $2n$ groups $D'_1, \dots, D'_n, D''_1, \dots, D''_n$ (where $n = \dim X$) and assume that the sets $S' = D'_1 \cap \cdots \cap D'_n$ and $S'' = D''_1 \cap \cdots \cap D''_n$ consist of special points only. The main result of this section is a certain reciprocity between the products of symbols over S' and S'' .

3.1 Toroidal pair

Here we recall the definition of a toroidal pair. A detailed treatment of toroidal pairs is given in [KKMS73] where they are called toroidal embeddings without self-intersections. We use Danilov’s terminology from [Dan78].

Let X be a normal n -dimensional variety over an algebraically closed field k . Let D be a closed subset of X , every irreducible component of which is a codimension-1 normal subvariety of X . We say that the pair (X, D) is *toroidal* if for every closed point $x \in X$ there exists an n -dimensional algebraic torus \mathbb{T} , an affine toric variety X_σ (corresponding to a rational convex n -dimensional cone σ), and a point x_0 in X_σ such that (X, D, x) is formally locally isomorphic to $(X_\sigma, X_\sigma \setminus \mathbb{T}, x_0)$. The latter means that there exists an isomorphism of the formal completions of the local rings

$$\widehat{\mathcal{O}}_{X,x} \cong \widehat{\mathcal{O}}_{X_\sigma, x_0},$$

such that the image of the ideal of D is mapped to the image of the ideal of $X_\sigma \setminus \mathbb{T}$. We call (X_σ, x_0) a *local model* of (X, D) at x .

Consider the n -form

$$\omega_0 = \frac{dt_1}{t_1} \wedge \cdots \wedge \frac{dt_n}{t_n},$$

where (t_1, \dots, t_n) are coordinates in \mathbb{T} . Automorphisms of \mathbb{T} correspond to monomial changes of coordinates

$$u_i = t_1^{q_{i1}} \cdots t_n^{q_{in}}, \quad 1 \leq i \leq n, \quad Q = (q_{ij}) \in GL(n, \mathbb{Z}). \quad (3)$$

We will write $u = t^Q$ to denote the monomial change of coordinates (3). Note that the form ω_0 is preserved under monomial changes of coordinates with $\det Q = 1$, and changes the sign when $\det Q = -1$. Therefore, ω_0 provides an analog of orientation on X_σ .

Furthermore, the choice of coordinates in \mathbb{T} defines an orientation of the space of characters $M \otimes \mathbb{R}$. Monomial changes of coordinates (3) preserve this orientation if and only if $\det Q = 1$. Therefore, the orientation of $M \otimes \mathbb{R}$ and hence of σ is uniquely determined by the form ω_0 .

We call $(X_\sigma, x_0, \omega_0)$ an *equipped local model* of (X, D) at x , assuming that the form ω_0 is fixed and the cone σ is oriented accordingly.

Let $D = \bigcup_{i \in I} E_i$ be the decomposition of D into irreducible components. The components of the sets $\bigcap_{i \in J} E_i \setminus \bigcup_{i \notin J} E_i$ (where $J \subset I$) are non-singular and define a stratification of X (see [KKMS73, p. 57]). In particular, $X \setminus D$ is non-singular. The components of $\bigcap_{i \in J} E_i$ are normal and are the closures of the strata. Furthermore, for each $x \in X$ the closures of strata which contain x correspond formally to the closures of the orbits in a local model (X_σ, x_0) at x .

We denote by $\text{St}_i(X)$ the set of all i -dimensional strata, and by $\overline{\text{St}}_i(X)$ the set of the closures of the i -dimensional strata. Note that if $x \in \text{St}_0(X)$ then in every local model (X_σ, x_0) at x the cone σ has an apex and x_0 is the closed orbit in X_σ .

In the next proposition we describe what coordinate transformations relate different local models at a point $x \in \text{St}_0(X)$.

PROPOSITION 3.1. *Let (X, D) be a toroidal pair, $x \in \text{St}_0(X)$. Then for any two local models (X_σ, x_0) and $(X_{\sigma'}, x'_0)$ at x , every isomorphism*

$$\pi : \widehat{\mathcal{O}}_{X_\sigma, x_0} \cong \widehat{\mathcal{O}}_{X_{\sigma'}, x'_0}$$

that maps the image of the ideal of $X_\sigma \setminus \mathbb{T}$ to the image of the ideal of $X_{\sigma'} \setminus \mathbb{T}'$ is induced by a change of coordinates of the form

$$u_i = \phi_i t_1^{q_{i1}} \cdots t_n^{q_{in}}, \quad \phi_i \in \widehat{\mathcal{O}}_{X_\sigma, x_0}^\times, \quad 1 \leq i \leq n, \quad Q = (q_{ij}) \in GL(n, \mathbb{Z}),$$

where t_1, \dots, t_n and u_1, \dots, u_n are coordinate functions on the tori \mathbb{T} and \mathbb{T}' , respectively.

Proof. Let $\Sigma(x)$ be the union of all strata Z whose closure \overline{Z} contains x . Denote by $M(x)$ the group of the Cartier divisors on $\Sigma(x)$, supported on $\Sigma(x) \cap D$, and by $M(x)_+$ the subsemigroup of effective divisors. For each local model (X_σ, x_0) at x , $M(x)$ is canonically isomorphic to the group of characters M of X_σ , and $M(x)_+$ is canonically isomorphic to the semigroup $\sigma \cap M$ (see [KKMS73, p. 61]). Therefore the semigroups $\sigma \cap M$ and $\sigma' \cap M'$ are isomorphic. In coordinates t_1, \dots, t_n and u_1, \dots, u_n , the isomorphism corresponds to a monomial transformation $u = t^Q$, for $Q \in GL(n, \mathbb{Z})$.

To describe all isomorphisms $\pi : \widehat{\mathcal{O}}_{X_\sigma, x_0} \cong \widehat{\mathcal{O}}_{X_{\sigma'}, x'_0}$ it suffices to describe all automorphisms α of $\widehat{\mathcal{O}}_{X_\sigma, x_0}$ that fix the orbits of X_σ . Let t_1, \dots, t_n be coordinates in \mathbb{T} . Then the ring $\widehat{\mathcal{O}}_{X_\sigma, x_0}$ can be identified with the ring of all formal power series in t_1, \dots, t_n supported in $\sigma \cap M$, where M is identified with \mathbb{Z}^n . Denote this ring by A . Let S be a multiplicative subset of A consisting of all elements ϕt^a , where $a \in \sigma \cap M$ and ϕ is an invertible element of A . Then for every automorphism α ,

we have $\alpha(S) \subseteq S$. Indeed, since α fixes the orbits of X_σ , it maps every ideal (t^a) to itself. Thus $\alpha(t^a) = \phi t^a$, for some invertible ϕ . Therefore, α induces an automorphism α_S of the localization A_S . Note that $t_1, \dots, t_n \in A_S$, since the elements of $\sigma \cap M$ generate M as a group. Therefore, for each i , $1 \leq i \leq n$, $\alpha_S(t_i) = \phi_i t_i$ for some invertible ϕ_i .

Conversely, every map $t_i \mapsto \phi_i t_i$, $1 \leq i \leq n$, $\phi_i \in A^\times$, defines an automorphism α of A , which fixes the orbits. Indeed, for every element $f \in A$, $f = \sum_a \lambda_a t^a$, $a \in \sigma \cap M$, put

$$\alpha(f) = \sum_{a \in \sigma \cap M} \lambda_a \phi^a t^a, \quad \phi^a = \phi_1^{a_1} \dots \phi_n^{a_n}. \quad (4)$$

Note that the coefficient of each monomial t^b of the series (4) is defined by finitely many series $\lambda_a \phi^a t^a$ (this is true since the cone σ has an apex). Therefore this is a well-defined power series. Since all the monomials in the series belong to the semigroup $\sigma \cap M$, the series defines an element of A . It is easy to check that α is in fact a homomorphism. Also it is clearly invertible. \square

3.2 Covering and combinatorial coefficients

DEFINITION 3.2. Let (X, D) be a toroidal pair. We say that (D_1, \dots, D_n) is a *reasonable covering* of D if $D = D_1 \cup \dots \cup D_n$, where each D_i is the union of some irreducible components of D , and $D_1 \cap \dots \cap D_n \subseteq \text{St}_0(X)$.

Let (D_1, \dots, D_n) be a reasonable covering of D . Consider an equipped local model $(X_\sigma, x_0, \omega_0)$ at a point $x \in \text{St}_0(X)$. It can be easily seen that the covering (D_1, \dots, D_n) defines a covering of the boundary of the cone σ in the sense of (2). This allows us to define the combinatorial coefficient of the covering (D_1, \dots, D_n) at each point $x \in \text{St}_0(X)$.

DEFINITION 3.3. The *combinatorial coefficient* at $x \in \text{St}_0(X)$ of the covering (D_1, \dots, D_n) is the combinatorial coefficient of the induced covering of σ in an equipped local model at x . We denote it by $c(x)$.

Remark 3.4 (Invariance). By Remark 2.5 the combinatorial coefficient is the same for any two equipped local models that correspond to an automorphism of \mathbb{T} that preserves the form ω_0 , and changes sign otherwise. Also it is skew-symmetric in D_1, \dots, D_n .

Now consider the stratification defined by the irreducible components of D (see § 3.1) and let F be a complete flag of stratum closures on X :

$$F : X_0 \subset X_1 \subset \dots \subset X_{n-1} \subset X, \quad X_i \in \overline{\text{St}}_i(X).$$

It corresponds to a complete flag of orbit closures in an equipped local model $(X_\sigma, x_0, \omega_0)$ of (X, D) at X_0 , hence to a complete flag F_σ of faces of σ .

DEFINITION 3.5. We say that the flag F is positive (respectively negative) and write $\text{sgn } F = 1$ (respectively $\text{sgn } F = -1$) if the induced flag F_σ of faces of σ is positive (respectively negative).

Like in the case of the combinatorial coefficient, the sign of the flag depends on the choice of the form ω_0 in an equipped local model.

DEFINITION 3.6. Let Z be a stratum. We say that the closure \overline{Z} has *signature* $\{i_1, \dots, i_k\}$ for $1 \leq i_l \leq n$ if and only if $Z \subseteq D_{i_1} \cap \dots \cap D_{i_k}$ and k is maximal.

The following proposition is the analog of the description of the combinatorial coefficient given in Theorem 2.4.

PROPOSITION 3.7. Let (X, D) be toroidal and (D_1, \dots, D_n) a reasonable covering of D . Then the combinatorial coefficient $c(x)$ of $x \in \text{St}_0(X)$ is equal to the number of all complete flags

$$x = X_0 \subset X_1 \subset \dots \subset X_{n-1} \subset X,$$

where $X_i \in \overline{\text{St}}_i(X)$ is a stratum closure of signature $\{i+1, \dots, n\}$, $0 \leq i \leq n-1$, counting signs.

3.3 Symbol of monomials

DEFINITION 3.8. Consider an ordered collection of $n+1$ monomials in n variables with coefficients in a field k :

$$c_i t^{a_i} = c_i t_1^{a_{i1}} \dots t_n^{a_{in}}, \quad c_i \in k^\times, \quad a_i = (a_{i1}, \dots, a_{in}) \in \mathbb{Z}^n, \quad 0 \leq i \leq n.$$

Let $A = (a_{ij}) \in M_{n+1,n}(\mathbb{Z})$ be the matrix whose rows are the vectors of exponents a_i . Then the symbol of $n+1$ monomials is the non-zero element of k defined by

$$[c_0 t^{a_0}, \dots, c_n t^{a_n}] = (-1)^B \prod_{i=0}^n c_i^{(-1)^i A_i},$$

where A_i is the determinant of the matrix obtained from A by eliminating its i th row, and

$$B = \sum_k \sum_{i < j} a_{ik} a_{jk} A_{ij}^k,$$

where A_{ij}^k is the determinant of the matrix obtained from A by eliminating its i th and j th rows and its k th column.

PROPOSITION 3.9. Let $f_i = c_i t^{a_i}$, $0 \leq i \leq n$, be monomials. The symbol has the following properties:

i) *Multiplicativity.* Suppose f_i is a product of two monomials $f_i = f'_i f''_i$. Then

$$[f_0, \dots, f'_i f''_i, \dots, f_n] = [f_0, \dots, f'_i, \dots, f_n] [f_0, \dots, f''_i, \dots, f_n].$$

ii) *Multiplicative skew-symmetry.*

$$[f_0, \dots, f_i, \dots, f_j, \dots, f_n] = [f_0, \dots, f_j, \dots, f_i, \dots, f_n]^{-1}.$$

iii) *Invariance.*

a) Let $u = t^Q$, $Q \in GL(n, \mathbb{Z})$ be a monomial change of coordinates. Then

$$[\bar{f}_0, \dots, \bar{f}_n] = [f_0, \dots, f_n]^{\det Q},$$

where $\bar{f}_i = c_i u^{a_i} = c_i t^{a_i Q}$ and $f_i = c_i t^{a_i}$, $0 \leq i \leq n$.

b) Let $s = \lambda t$ be a translation, i.e. $s_j = \lambda_j t_j$, $\lambda_j \in k^\times$, $1 \leq j \leq n$. Then

$$[f'_0, \dots, f'_n] = [f_0, \dots, f_n],$$

where $f'_i = c_i s^{a_i} = c_i \lambda^{a_i} t^{a_i}$ and $f_i = c_i t^{a_i}$, $0 \leq i \leq n$.

Proof. Modulo the sign $(-1)^B$ all the properties follow easily from the properties of the determinant.

To take care of the sign we give an invariant description of B , following [Kho99]. Consider B as a $\mathbb{Z}/2\mathbb{Z}$ -valued function of the rows a_0, \dots, a_n of the matrix A . It is easy to see that $B = B(a_0, \dots, a_n)$ is multilinear and its value on each collection of $n+1$ standard vectors $(e_{i_0}, \dots, e_{i_n})$ is 0 if more than two of the vectors e_{i_0}, \dots, e_{i_n} coincide; and 1 otherwise.

Now define the function $B' = B'(a_0, \dots, a_n)$ to be 0 if the rank of (a_0, \dots, a_n) is less than n ; and $\lambda_0 + \dots + \lambda_n + 1 \pmod{2}$ if the vectors a_0, \dots, a_n satisfy a (unique) non-trivial relation $\lambda_0 a_0 + \dots + \lambda_n a_n = 0$. The function B' is multilinear and on each collection $(e_{i_0}, \dots, e_{i_n})$ the functions B' and B take the same value. Therefore $B = B'$; in particular, B is symmetric and invariant under non-degenerate transformations. \square

3.4 Toroidal symbol

Let (X, D) be toroidal. Let $k(X, D)$ denote the set of rational functions on X whose divisor lies in D .

Consider a zero-dimensional stratum $x \in \text{St}_0(X)$. Then the image of $f \in k(X, D)$ in an equipped local model $(X_\sigma, x_0, \omega_0)$ at x is the product of a monomial ct^a and a regular invertible function $\phi \in \hat{\mathcal{O}}_{X_\sigma, x_0}^\times$ with $\phi(x_0) = 1$. We call this monomial the *leading monomial of f at x* . The leading monomial is defined up to monomial transformations.

DEFINITION 3.10. Let (X, D) be toroidal and $x \in \text{St}_0(X)$ a zero-dimensional stratum. Define the *toroidal symbol* $[f_0, \dots, f_n]_x$ at x of $f_0, \dots, f_n \in k(X, D)$ to be the symbol of the leading monomials of f_0, \dots, f_n at x .

Remark 3.11 (Invariance). Let $(X_{\sigma'}, x'_0, \omega'_0)$ and $(X_{\sigma''}, x''_0, \omega''_0)$ be two equipped local models at x . Let f' and f'' be the images of $f \in k(X, D)$ in the two equipped local models. Then, according to Proposition 3.1, the leading monomials of f' and f'' are related by a composition of a monomial transformation and a translation: $t \mapsto \lambda t^Q$. Therefore, by Proposition 3.9 the toroidal symbol is the same for the two equipped local models if $\det Q = 1$, and is reciprocal otherwise.

By Proposition 3.9 the toroidal symbol is multiplicative and multiplicatively skew-symmetric in f_0, \dots, f_n .

Now we will give a relation between the toroidal symbol and Parshin's tame symbol at a complete flag of stratum closures on X .

PROPOSITION 3.12. Let (X, D) be toroidal. Consider $n+1$ rational functions $f_0, \dots, f_n \in k(X, D)$. Then, for any complete flag $F : X_0 \subset X_1 \subset \dots \subset X_{n-1} \subset X$ of stratum closures on X , we have

$$\langle f_0, \dots, f_n \rangle_F = [f_0, \dots, f_n]_{X_0}^{\text{sgn } F},$$

where $\langle f_0, \dots, f_n \rangle_F$ denotes Parshin's tame symbol at the flag F .

(Note that the number $[f_0, \dots, f_n]_{X_0}^{\text{sgn } F}$ is already independent of the choice of ω_0 in an equipped local model.)

Proof. Since the definitions of the toroidal symbol and Parshin's tame symbol are local, we can pass to an equipped local model $(X_\sigma, x_0, \omega_0)$ at X_0 and assume that F is a complete flag of orbit closures in X_σ .

Let F_σ be the complete flag of faces of σ corresponding to F ,

$$F_\sigma : \quad 0 = \sigma_0 \subset \sigma_1 \subset \dots \subset \sigma_{n-1} \subset \sigma_n = \sigma.$$

Fix coordinates (t_1, \dots, t_n) in \mathbb{T} , $M \cong \mathbb{Z}^n$. Inside each σ_i , $1 \leq i \leq n$, choose a lattice point $q_i \in \mathbb{Z}^n$ at lattice distance 1 from σ_{i-1} . Let $u_i = t^{q_i}$ be a monomial change of coordinates in \mathbb{T} . Then the rational functions $u_i = t^{q_i}$ give a system of local parameters at F (see Appendix). Therefore by Proposition 3.9 and Remark A.2 of the Appendix,

$$\langle f_0, \dots, f_n \rangle_F = [c_0 u^{k_0}, \dots, c_n u^{k_n}] = [c_0 t^{k_0}, \dots, c_n t^{k_n}]^{\det Q},$$

where $Q = (q_1, \dots, q_n)$ and $c_i t^{k_i}$ is the leading monomial of f_i . It remains to note that $\det Q = \text{sgn } F$. \square

COROLLARY 3.13. Let (X, D) be toroidal and (D_1, \dots, D_n) be a reasonable covering of D . For $x \in \text{St}_0(X)$ let $\mathcal{F}(x)$ be the set of all complete flags

$$x = X_0 \subset X_1 \subset \dots \subset X_{n-1} \subset X,$$

where $X_i \in \overline{\text{St}}_i(X)$ is a stratum closure of signature $\{i+1, \dots, n\}$, $0 \leq i \leq n-1$.

Then for any $n + 1$ rational functions $f_0, \dots, f_n \in k(X, D)$ we have

$$[f_0, \dots, f_n]_x^{c(x)} = \prod_{F \in \mathcal{F}(x)} \langle f_0, \dots, f_n \rangle_F,$$

where we assume that the product is 1 if $\mathcal{F}(x)$ is empty.

Proof. This follows from Propositions 3.7 and 3.12. \square

Remark 3.14. Note that for a toroidal pair (X, D) all Parshin's reciprocity laws (Theorem A.4) for $i > 0$ follow from Proposition 3.12. Indeed, consider a complete flag $F : X_0 \subset \dots \subset X_i \subset \dots \subset X_n$ of irreducible subvarieties of X . We can pass to a local model at X_0 and assume that X is an affine toric variety and $D = X \setminus \mathbb{T}$. Then for any $n + 1$ rational functions $f_0, \dots, f_n \in k(X, D)$ the symbol $\langle f_0, \dots, f_n \rangle_F$ is trivial unless F is a flag of orbit closures on X . But if we fix all X_j , $j \neq i$, and vary X_i , there are only two such flags F (since for any face τ of a polyhedral cone there are only two codimension-1 faces of τ that contain a fixed codimension-2 face of τ), and the signs of these two flags are opposite. Now we can apply Proposition 3.12.

3.5 Main theorem

THEOREM 3.15. *Let X be a complete normal n -dimensional variety over an algebraically closed field k , and D a closed subset of X such that the pair (X, D) is toroidal.*

Let (D_1, \dots, D_n) be a reasonable covering of D such that each D_i is a disjoint union of two closed subsets of pure codimension 1:

$$D = D_1 \cup \dots \cup D_n, \quad D_1 \cap \dots \cap D_n \subseteq \text{St}_0(X), \quad D_i = D'_i \sqcup D''_i, \quad 1 \leq i \leq n. \quad (5)$$

We get 2^n disjoint finite closed subsets of X :

$$S_k = G_1 \cap \dots \cap G_n, \quad \text{where } G_i = D'_i \text{ or } D''_i, \quad 1 \leq i \leq n, \quad 1 \leq k \leq 2^n.$$

Then for any $n + 1$ rational functions $f_0, \dots, f_n \in k(X, D)$, the following 2^n numbers are equal:

$$\left(\prod_{x \in S_1} [f_0, \dots, f_n]_x^{c(x)} \right)^{(-1)^{|S_1|}} = \dots = \left(\prod_{x \in S_{2^n}} [f_0, \dots, f_n]_x^{c(x)} \right)^{(-1)^{|S_{2^n}|}}, \quad (6)$$

where $[f_0, \dots, f_n]_x$ is the toroidal symbol of f_0, \dots, f_n at x , $c(x)$ is the combinatorial coefficient at x , and $|S_k|$ is the number of D''_i in the definition of S_k .

Proof. Because of the symmetry it is sufficient to prove the equality for any two sets

$$S_1 = D'_1 \cap \dots \cap D'_i \cap \dots \cap D'_n \quad \text{and} \quad S_2 = D'_1 \cap \dots \cap D''_i \cap \dots \cap D'_n.$$

Since the number $[f_0, \dots, f_n]_x^{c(x)}$ is multiplicatively skew-symmetric in D_1, \dots, D_n (see Remark 3.4) we may assume that $i = 1$, so

$$S_1 = D'_1 \cap D'_2 \cap \dots \cap D'_n \quad \text{and} \quad S_2 = D''_1 \cap D'_2 \cap \dots \cap D'_n.$$

We have to show that

$$\prod_{x \in S_1 \cup S_2} [f_0, \dots, f_n]_x^{c(x)} = 1.$$

Let Σ be the union of all stratum closures $Y \in \overline{\text{St}}_1(X)$ with signature $\{2', \dots, n'\}$. It follows from Corollary 3.13 that if $x \in S_1 \cup S_2$ does not lie on any component of Σ then $[f_0, \dots, f_n]_x^{c(x)} = 1$. On the other hand, by (5) the signature of every point $x \in \text{St}_0(X) \cap \Sigma$ is either

$$\{2', \dots, n'\} \quad \text{or} \quad \{1', 2', \dots, n'\} \quad \text{or} \quad \{1'', 2', \dots, n'\}.$$

In the first case $[f_0, \dots, f_n]_x^{c(x)} = 1$, again by Corollary 3.13. In the second case $x \in S_1$; and in the third $x \in S_2$. Therefore, we have

$$\prod_{x \in S_1 \cup S_2} [f_0, \dots, f_n]_x^{c(x)} = \prod_{x \in \text{St}_0(X) \cap \Sigma} [f_0, \dots, f_n]_x^{c(x)}. \quad (7)$$

Now consider a component Y of Σ , and a closed point $y \in Y$. Let $\mathcal{F}(y, Y)$ denote the set of all complete flags

$$y \subset Y \subset X_2 \subset \dots \subset X_{n-1} \subset X,$$

where $X_i \in \overline{\text{St}}_i(X)$ has signature $\{(i+1)', \dots, n'\}$, $2 \leq i \leq n-1$. Denote

$$\langle f_0, \dots, f_n \rangle_{y, Y} = \prod_{F \in \mathcal{F}(y, Y)} \langle f_0, \dots, f_n \rangle_F,$$

and assume that $\langle f_0, \dots, f_n \rangle_{y, Y} = 1$ if $\mathcal{F}(y, Y)$ is empty. Then by Corollary 3.13 for each $x \in \text{St}_0(X) \cap \Sigma$ we have

$$[f_0, \dots, f_n]_x^{c(x)} = \prod_{F \in \mathcal{F}(x)} \langle f_0, \dots, f_n \rangle_F = \prod_{Y \ni x} \langle f_0, \dots, f_n \rangle_{x, Y}, \quad (8)$$

where the product on the right-hand side runs over all components Y of Σ containing x .

On the other hand, by the first Parshin's reciprocity law (Theorem A.4 for $i = 0$)

$$\prod_{y \in Y} \langle f_0, \dots, f_n \rangle_{y \subset Y \subset X_2 \subset \dots \subset X_{n-1} \subset X} = 1,$$

where the product is taken over all points $y \in Y$. Thus

$$\prod_{y \in Y} \langle f_0, \dots, f_n \rangle_{y, Y} = 1.$$

Note that $\langle f_0, \dots, f_n \rangle_{y, Y}$ is trivial for all points y not lying in $\text{St}_0(X)$, so we can assume that $y \in \text{St}_0(X) \cap Y$. We have

$$\prod_{y \in \text{St}_0(X) \cap Y} \langle f_0, \dots, f_n \rangle_{y, Y} = 1. \quad (9)$$

Combining (7), (8) and (9) we get

$$\begin{aligned} \prod_{x \in S_1 \cup S_2} [f_0, \dots, f_n]_x^{c(x)} &= \prod_{x \in \text{St}_0(X) \cap \Sigma} [f_0, \dots, f_n]_x^{c(x)} \\ &= \prod_{x \in \text{St}_0(X) \cap \Sigma} \prod_{Y \ni x} \langle f_0, \dots, f_n \rangle_{x, Y} \\ &= \prod_{Y \subset \Sigma} \prod_{x \in \text{St}_0(X) \cap Y} \langle f_0, \dots, f_n \rangle_{x, Y} = 1. \end{aligned} \quad \square$$

4. Toroidal residue

Let (X, D) be a toroidal pair, as before. At each point $x \in \text{St}_0(X)$ we define the residue $\text{res}_x^{\mathbb{T}} \omega$ of a rational n -form ω on X which is regular in $X \setminus D$. Then we prove an additive analog of Theorem 3.15.

4.1 Toroidal residue

First we will define the toroidal residue for a local model $(X_\sigma, x_0, \omega_0)$ at a point $x \in \text{St}_0(X)$. As before X_σ is an n -dimensional affine toric variety, x_0 is the closed orbit, and $\omega_0 = dt_1/t_1 \wedge \dots \wedge dt_n/t_n$, where (t_1, \dots, t_n) is a coordinate system in \mathbb{T} .

Let $A = \hat{\mathcal{O}}_{X_\sigma, x_0}$ be the completion of the local ring of x_0 on X_σ , and $B = A_S$ the localization of A by the multiplicative subgroup S of all monomials. We consider the B -algebra Ω_B^n of differential n -forms that are regular in \mathbb{T} , and the A -algebra Ω_A^n of regular differential n -forms.

By fixing coordinates (t_1, \dots, t_n) we can identify every element $f \in B$ with a formal power series

$$f = t^b \sum_{a \in \sigma \cap \mathbb{Z}^n} \lambda_a t^a, \quad b \in \mathbb{Z}^n.$$

Let $\omega \in \Omega_B^n$. Then we can write $\omega = f dt_1/t_1 \wedge \dots \wedge dt_n/t_n$, for some $f \in B$.

DEFINITION 4.1. The *toroidal residue* of a differential n -form $\omega \in \Omega_B^n$ is the constant term λ_{-b} in the formal power series of f . We denote it by $\text{res}^\mathbb{T} \omega$.

We have the following properties of the toroidal residue.

PROPOSITION 4.2. Consider a differential n -form $\omega \in \Omega_B^n$. Then the following hold.

- i) If ω is exact then $\text{res}^\mathbb{T} \omega = 0$.
- ii) If $\omega \in \Omega_A^n$ then $\text{res}^\mathbb{T} \omega = 0$.
- iii) For any $s_1, \dots, s_n \in B^\times$, $\text{res}^\mathbb{T} ds_1/s_1^{m_1} \wedge \dots \wedge ds_n/s_n^{m_n} = 0$, unless all $m_i = 1$.
- iv) The toroidal residue is independent of the choice of coordinates (t_1, \dots, t_n) .
- v) The toroidal residue is invariant under monomial transformations $t \mapsto t^Q$, $Q \in GL(n, \mathbb{Z})$, up to a factor $\det Q$.

Proof. i) Let $\omega = dw$. We can assume that $w = g dt_1 \wedge \dots \wedge \widehat{dt_i} \wedge \dots \wedge dt_n$. Then

$$\omega = (-1)^{i-1} \frac{\partial g}{\partial t_i} t_1 \dots t_n \frac{dt_1}{t_1} \wedge \dots \wedge \frac{dt_n}{t_n}.$$

Suppose $g = \sum_a \lambda_a t^a$. Then

$$(-1)^{i-1} \frac{\partial g}{\partial t_i} t_1 \dots t_n = \sum_a a_i \lambda_a \frac{t^a}{t_i} t_1 \dots t_n, \quad a = (a_1, \dots, a_n).$$

Clearly the constant term of the last series is zero.

ii) Let $u_i = t^{a_i}$, $a_i \in \sigma \cap \mathbb{Z}^n$, be n regular functions, such that du_1, \dots, du_n are linearly independent. Then for every $\omega \in \Omega_A^n$

$$\omega = f du_1 \wedge \dots \wedge du_n = f dt^{a_1} \wedge \dots \wedge dt^{a_n} = f t^{a_1 + \dots + a_n} J \frac{dt_1}{t_1} \wedge \dots \wedge \frac{dt_n}{t_n},$$

where $f \in A$ and $J = \det(a_1, \dots, a_n)$. Clearly, the constant term of $f t^{a_1 + \dots + a_n} J$ is zero.

iii) First assume that $\text{char}(k) = 0$. Suppose $m_i \neq 1$. Then

$$\frac{ds_1}{s_1^{m_1}} \wedge \dots \wedge \frac{ds_n}{s_n^{m_n}} = d \left(\frac{(-1)^{i-1}}{1 - m_i} s_i^{1-m_i} \frac{ds_1}{s_1^{m_1}} \wedge \dots \wedge \frac{\widehat{ds_i}}{s_i^{m_i}} \wedge \dots \wedge \frac{ds_n}{s_n^{m_n}} \right),$$

and the statement follows from part i. In the case of an arbitrary characteristic, note that the toroidal residue is a polynomial function in finitely many coefficients of the series $s_1, \dots, s_n \in B^\times$. This function is independent of the characteristic and vanishes when the characteristic is zero. Therefore it is identically zero.

iv) Let (s_1, \dots, s_n) be another coordinate system in \mathbb{T} . Then $s_i = \phi_i t_i$, where $\phi_i \in A^\times$. Consider $\omega \in \Omega_B^n$ and write

$$\omega = f \frac{ds_1}{s_1} \wedge \dots \wedge \frac{ds_n}{s_n}, \quad f = \sum_a \lambda_a s^a, \quad a \in \mathbb{Z}^n.$$

Then the residue of ω with respect to (s_1, \dots, s_n) is $\text{res}_{(s_1, \dots, s_n)}^{\mathbb{T}} \omega = \lambda_0$. On the other hand, by part iii the residue of ω with respect to (t_1, \dots, t_n) equals

$$\text{res}_{(t_1, \dots, t_n)}^{\mathbb{T}} \omega = \text{res}_{(t_1, \dots, t_n)}^{\mathbb{T}} \lambda_0 \frac{ds_1}{s_1} \wedge \dots \wedge \frac{ds_n}{s_n}. \quad (10)$$

Now, for each $1 \leq i \leq n$,

$$\frac{ds_i}{s_i} = \frac{d\phi_i}{\phi_i} + \frac{dt_i}{t_i}.$$

Substituting into (10) and expanding we get

$$\text{res}_{(t_1, \dots, t_n)}^{\mathbb{T}} \omega = \text{res}_{(t_1, \dots, t_n)}^{\mathbb{T}} \lambda_0 \frac{dt_1}{t_1} \wedge \dots \wedge \frac{dt_n}{t_n} + \sum_i \text{res}_{(t_1, \dots, t_n)}^{\mathbb{T}} \omega_i,$$

where in each ω_i at least one of the dt_j/t_j is replaced by $d\phi_j/\phi_j$. It is easy to check that the residue of every ω_i is zero.

v) This follows from the fact that if $u_i = t^{q_i}$, $q_i \in \mathbb{Z}^n$, then

$$\frac{du_1}{u_1} \wedge \dots \wedge \frac{du_n}{u_n} = \det Q \frac{dt_1}{t_1} \wedge \dots \wedge \frac{dt_n}{t_n},$$

and from the observation that monomial transformations do not change the constant term of a series. \square

Remark 4.3. The proof of parts iii and iv is similar to the one given in [FP04] for Parshin's residue.

Now let (X, D) be toroidal. Denote by $\Omega^n(X, D)$ the set of all rational n -forms on X that are regular in $X \setminus D$.

DEFINITION 4.4. The *toroidal residue* of a rational n -form $\omega \in \Omega^n(X, D)$ at a point $x \in \text{St}_0(X)$ is the toroidal residue of its image in a local model at x . We denote it by $\text{res}_x^{\mathbb{T}} \omega$.

Remark 4.5 (Invariance). As follows from Propositions 3.1 and 4.2, the toroidal residue is the same for any two equipped local models that are related by an isomorphism preserving the form ω_0 , and changes sign otherwise.

The relation between the toroidal residue and Parshin's residue is similar to the one between the toroidal symbol and Parshin's tame symbol.

PROPOSITION 4.6. Let (X, D) be toroidal. Consider a rational n -form $\omega \in \Omega^n(X, D)$. Then for any complete flag $F : X_0 \subset X_1 \subset \dots \subset X_{n-1} \subset X$ of stratum closures on X we have

$$\text{res}_F \omega = \text{sgn } F \text{res}_{X_0}^{\mathbb{T}} \omega,$$

where $\text{res}_F \omega$ denotes Parshin's residue at the flag F .

The number $\text{sgn } F \text{res}_x^{\mathbb{T}} \omega$ is independent of the choice of the form ω_0 in an equipped local model. Consequently, if (D_1, \dots, D_n) is a reasonable covering of D then at each $x \in \text{St}_0(X)$ the number $c(x) \text{res}_x^{\mathbb{T}} \omega$ is also well defined.

The following is the additive analog of Corollary 3.13.

COROLLARY 4.7. Let (X, D) be toroidal and (D_1, \dots, D_n) be a reasonable covering of D . For $x \in \text{St}_0(X)$ let $\mathcal{F}(x)$ be the set of all complete flags

$$x = X_0 \subset X_1 \subset \dots \subset X_{n-1} \subset X,$$

where $X_i \subset \overline{\text{St}_i(X)}$ is a stratum closure of signature $\{i+1, \dots, n\}$, $0 \leq i \leq n-1$.

Then for any $\omega \in \Omega^n(X, D)$ we have

$$c(x) \operatorname{res}_x^{\mathbb{T}} \omega = \sum_{F \in \mathcal{F}(x)} \operatorname{res}_F \omega,$$

where we assume that the sum is 0 if $\mathcal{F}(x)$ is empty.

4.2 Main theorem

THEOREM 4.8. *Let X be a complete normal n -dimensional variety over an algebraically closed field k , and D a closed subset of X such that the pair (X, D) is toroidal.*

Let (D_1, \dots, D_n) be a reasonable covering of D such that each D_i is a disjoint union of two closed subsets of pure codimension 1:

$$D = D_1 \cup \dots \cup D_n, \quad D_1 \cap \dots \cap D_n \subset \operatorname{St}_0(X), \quad D_i = D'_i \sqcup D''_i, \quad 1 \leq i \leq n. \quad (11)$$

We get 2^n disjoint finite closed subsets of X :

$$S_k = G_1 \cap \dots \cap G_n, \quad \text{where } G_i = D'_i \text{ or } D''_i, \quad 1 \leq i \leq n, \quad 1 \leq k \leq 2^n.$$

Then for any rational n -form $\omega \in \Omega^n(X, D)$, the following 2^n numbers are equal:

$$(-1)^{|S_1|} \sum_{x \in S_1} c(x) \operatorname{res}_x^{\mathbb{T}} \omega = \dots = (-1)^{|S_{2^n}|} \sum_{x \in S_{2^n}} c(x) \operatorname{res}_x^{\mathbb{T}} \omega,$$

where $\operatorname{res}_x^{\mathbb{T}} \omega$ denotes the toroidal residue of ω at x , $c(x)$ is the combinatorial coefficient at x , and $|S_k|$ is the number of D''_i in the definition of S_k .

Proof. The proof repeats the arguments of the proof of Theorem 3.15. □

5. Application to systems of equations in the torus

In this section we apply our main results on the toroidal symbol and residue to prove the product of roots formula and the sum of values formula (Theorems A and B in the Introduction).

Recall that a *Laurent polynomial* is a finite linear combination of monomials with integer exponent vectors and coefficients in k :

$$f(t) = \sum_{m \in \mathbb{Z}^n} \lambda_m t^m, \quad t^m = t_1^{m_1} \dots t_n^{m_n}, \quad \lambda_m \in k.$$

The convex hull of those lattice points $m \in \mathbb{Z}^n$ for which $\lambda_m \neq 0$ is called the *Newton polytope* of f . The value of $f(t)$ is defined for all t in the algebraic n -torus $\mathbb{T} = (k^\times)^n$.

Consider a system of n Laurent polynomial equations in \mathbb{T} :

$$f_1(t) = \dots = f_n(t) = 0, \quad t \in \mathbb{T}. \quad (12)$$

Let Δ_i be the Newton polytope of f_i . We assume that none of the f_i is a monomial, hence, none of Δ_i is a point.

Every linear functional w on \mathbb{R}^n defines a collection of faces $\Delta_1^w, \dots, \Delta_n^w$ of the Newton polytopes such that the restriction of w on Δ_i attains its maximum precisely at Δ_i^w . The polynomial

$$f_i^w(t) = \sum_{m \in \Delta_i^w \cap \mathbb{Z}^n} \lambda_m t^m$$

is called the *initial form* of f_i with respect to w . According to Bernstein's theorem [Ber75] the number of solutions to the system (12) is finite (and equals the mixed volume of $\Delta_1, \dots, \Delta_n$) if and only if for every $w \neq 0$ the system $f_1^w(t) = \dots = f_n^w(t) = 0$ is inconsistent.

DEFINITION 5.1. A collection of polytopes $\Delta_1, \dots, \Delta_n$ is called *developed* if none of them is a point and for each $w \neq 0$ at least one of the faces $\Delta_1^w, \dots, \Delta_n^w$ is a vertex.

By the above any system with developed collection of Newton polytopes has finitely many solutions. We will call them the *roots* of the system. Every root x of the system has a multiplicity $\mu(x)$.

Let Δ be the Minkowski sum of $\Delta_1, \dots, \Delta_n$. Then every face $\Gamma \subset \Delta$ has a unique decomposition as a sum of faces

$$\Gamma = \Gamma_1 + \dots + \Gamma_n, \quad \text{where } \Gamma_i \subset \Delta_i, \quad i = 1, \dots, n. \quad (13)$$

If the collection $\Delta_1, \dots, \Delta_n$ is developed then Δ has dimension n and in the decomposition of every proper face of Δ at least one summand is a vertex. In this case, for each vertex A of Δ , the combinatorial coefficient $c(A)$ is defined.

DEFINITION 5.2. Let σ_A be the cone with apex A generated by the facets of Δ that contain A . Then the boundary of σ_A is covered by the closed sets D_1, \dots, D_n , where D_i is the union of all facets of σ_A that correspond to the facets of Δ whose i th summand in the decomposition (13) is a vertex. The combinatorial coefficient of this covering is called the *combinatorial coefficient* $c(A)$ of the vertex $A \in \Delta$.

5.1 Product of roots formula

Consider a system of n Laurent polynomials with developed collection of Newton polytopes. The product of the roots counting multiplicities is a point in \mathbb{T} , which we denote by ρ . To locate ρ it is enough to find the product of the values of t_i over the roots of the system, for each $1 \leq i \leq n$. More generally, we will find the product of the values of any Laurent monomial ct^m , for $c \in k^\times$, $m \in \mathbb{Z}^n$, over the roots of the system.

DEFINITION 5.3 [Kho99]. The symbol of f_1, \dots, f_n and a Laurent monomial $f_0 = ct^m$ at a vertex $A \in \Delta$ is the symbol of $n+1$ monomials $[ct^m, f_1(A_1)t^{A_1}, \dots, f_n(A_n)t^{A_n}]$, where $A = A_1 + \dots + A_n$ is the decomposition of A , and $f_i(A_i)$ is the coefficient of t^{A_i} in f_i . We denote it by $[f_0, \dots, f_n]_A$.

The following theorem was proved by Khovanskii [Kho99] in the complex case. Our proof uses the result of Theorem 3.15 and works over an arbitrary algebraically closed field k .

THEOREM 5.4. Suppose the collection of the Newton polytopes $\Delta_1, \dots, \Delta_n$ of the system (12) is developed. Then the product of the values of a Laurent monomial f_0 over the roots of the system is given by

$$\prod_{f_1(x)=\dots=f_n(x)=0} f_0(x)^{\mu(x)} = \prod_A [f_0, \dots, f_n]_A^{(-1)^n c(A)}, \quad (14)$$

where the right-hand product is taken over all vertices A of the polytope $\Delta = \Delta_1 + \dots + \Delta_n$ and $c(A)$ is the combinatorial coefficient at A .

Proof. First let us notice that it is sufficient to prove the theorem for a generic system with given Newton polytopes. There is an open set U in the space of the coefficients of the system where the number of roots counting multiplicities is constant. The left-hand side of (14), being symmetric in the roots of the system, is a rational function in the coefficients of the system. On the other hand, the product of the symbols $[f_0, \dots, f_n]_A$ is also a rational function of the coefficients of the system. Suppose we proved the formula for almost all systems in U . Then the two rational functions coincide on an open algebraic subset $W \subset U$, and thus coincide everywhere in U .

Consider the complete toric variety X associated with the Minkowski sum Δ (e.g. see [Ful93]). Let $D = X \setminus \mathbb{T}$ be the invariant divisor on X . Denote by Z_i the closure of the zero locus $f_i = 0$ in X ,

and let $Z = Z_1 \cup \dots \cup Z_n$. If the system is generic, the components of Z intersect transversally and the intersection of each component of Z with D is also transversal. In this case the pair $(X, D \cup Z)$ is toroidal.

Now we will define a covering of $D \cup Z$. Each irreducible component of D corresponds to a facet of Δ . Recall that each facet Γ of Δ has a unique decomposition into the sum of faces (13). Denote by D_i the union of all components that correspond to those facets whose i th summand is a vertex. Since the collection $\Delta_1, \dots, \Delta_n$ is developed, the sets D_1, \dots, D_n define a covering of D .

Consider a covering $D \cup Z = (D_1 \cup Z_1) \cup \dots \cup (D_n \cup Z_n)$. Notice that every intersection point in $(D_1 \cup Z_1) \cap \dots \cap (D_n \cup Z_n)$ is either a fixed orbit or a transversal intersection of some components of Z and D . Thus the covering is reasonable (see § 3.2). By definition, the components of D_i correspond to facets of Δ whose i th summand is a vertex, i.e. the corresponding initial forms of f_i are monomials. This implies that $D_i \cap Z_i = \emptyset$. Now applying Theorem 3.15 we obtain

$$\prod_{x \in Z_1 \cap \dots \cap Z_n} [f_0, \dots, f_n]_x^{c(x)} = \prod_{x \in D_1 \cap \dots \cap D_n} [f_0, \dots, f_n]_x^{(-1)^n c(x)}.$$

It remains to notice that for each transversal intersection x of components of Z the combinatorial coefficient $c(x) = 1$ and $[f_0, \dots, f_n]_x = f_0(x)^{\mu(x)}$ (see Example 1 in the Appendix). Also for a point $x \in D_1 \cap \dots \cap D_n$ the toroidal symbol $[f_0, \dots, f_n]_x$ coincides with $[f_0, \dots, f_n]_A$, where A is the corresponding vertex of Δ , and $c(x) = c(A)$, by definition. \square

5.2 Sum of values formula

We recall the definition of a Laurent series at a vertex and the residue at a vertex from [GK96]. Let f be a Laurent polynomial with Newton polytope $\Delta(f)$, A a vertex of $\Delta(f)$, and $f(A)$ the coefficient of t^A in f . Since the constant term of the Laurent polynomial $\tilde{f} = f/(f(A)t^A)$ equals 1, we get a well-defined power series

$$\frac{1}{\tilde{f}} = 1 + (1 - \tilde{f}) + (1 - \tilde{f})^2 + \dots. \quad (15)$$

DEFINITION 5.5. Let g be a Laurent polynomial. The formal product of the series (15) and the Laurent polynomial $g/(f(A)t^A)$ is called the *Laurent series of g/f at the vertex $A \in \Delta(f)$* .

DEFINITION 5.6. The *residue at a vertex $A \in \Delta(f)$* of a rational n -form

$$\omega = \frac{g}{f} \frac{dt_1}{t_1} \wedge \dots \wedge \frac{dt_n}{t_n}$$

is the constant term of the Laurent series of g/f at A . We denote it by $\text{res}_A \omega$.

The following theorem was proved by Gel'fond and Khovanskii [GK02] in the case when $k = \mathbb{C}$. We prove it for any algebraically closed field k using Theorem 4.8.

THEOREM 5.7. Suppose the collection of the Newton polytopes $\Delta_1, \dots, \Delta_n$ of the system (12) is developed. Then the sum of the values of a Laurent polynomial f_0 over the roots of the system counting multiplicities is given by

$$\sum_{f_1(x)=\dots=f_n(x)=0} \mu(x) f_0(x) = (-1)^n \sum_A c(A) \text{res}_A \left(\frac{f_0 J}{f} \frac{dt_1}{t_1} \wedge \dots \wedge \frac{dt_n}{t_n} \right),$$

where the sum on the right is taken over the vertices A of $\Delta = \Delta_1 + \dots + \Delta_n$, $J = \det(t_j \partial f_i / \partial t_j)$ for $1 \leq i, j \leq n$, $f = f_1 \dots f_n$, and $c(A)$ is the combinatorial coefficient at A .

Proof. As in the proof of Theorem 5.4 it is enough to consider the case of a generic system with given Newton polytopes, since the sum of the values of a Laurent polynomial over the roots of the system is a rational function of the coefficients of the system.

As before, X is the complete toric variety associated with the Minkowski sum, D the invariant divisor on X , Z_i the closure of the zero locus $f_i = 0$ in X , and $D \cup Z = (D_1 \cup Z_1) \cup \cdots \cup (D_n \cup Z_n)$ a reasonable covering that satisfies $D_i \cap Z_i = \emptyset$. Applying Theorem 4.8 to the form

$$\omega = f_0 \frac{df_1}{f_1} \wedge \cdots \wedge \frac{df_n}{f_n}$$

we get

$$\sum_{x \in Z_1 \cap \cdots \cap Z_n} c(x) \operatorname{res}_x^{\mathbb{T}} \omega = (-1)^n \sum_{x \in D_1 \cap \cdots \cap D_n} c(x) \operatorname{res}_x^{\mathbb{T}} \omega.$$

For each transversal intersection x of components of Z the combinatorial coefficient $c(x) = 1$ and $\operatorname{res}_x^{\mathbb{T}} \omega = \mu(x) f_0(x)$ (see Example 2 in the Appendix). Also for a point $x \in D_1 \cap \cdots \cap D_n$, we have $c(x) = c(A)$ and

$$\operatorname{res}_x^{\mathbb{T}} \omega = \operatorname{res}_A \left(\frac{f_0 J}{f} \frac{dt_1}{t_1} \wedge \cdots \wedge \frac{dt_n}{t_n} \right),$$

where $A \in \Delta$ is the vertex corresponding to the fixed orbit x . Therefore, we have obtained the required equality. \square

ACKNOWLEDGEMENTS

The results of the present paper constitute a part of the author's PhD thesis [Sop02]. I am grateful to Askold Khovanskii for stating the problem, his constant support and numerous discussions.

Appendix. Parshin's reciprocity laws

Here we recall the definition of Parshin's tame symbol and residue for an arbitrary algebraic variety X over an algebraically closed field, and formulate Parshin's reciprocity laws.

A.1 Parshin's tame symbol

Let X be a complete algebraic variety over an algebraically closed field k .

Consider a complete flag of irreducible subvarieties of X :

$$F : X_0 \subset X_1 \subset \cdots \subset X_{n-1} \subset X_n = X. \quad (\text{A1})$$

We will assume that all X_i are normal. The general case can be reduced to this one by considering normalization (for details see [Par83] or [Sop02]). Note also that this assumption holds for complete flags of stratum closures on toroidal pairs.

Given a flag F as in (A1) define a *system of local parameters at F* , $(u_1, \dots, u_n)_F \in k(X)^n$, as follows. There exists an open subset of X_n where the codimension-1 subvariety X_{n-1} has a local equation u_n . In general, for every $i = 0, \dots, n-1$, let u_{n-i} be a local equation (in some open subset) of the codimension-1 subvariety $X_{n-i-1} \subset X_{n-i}$.

Next, for every rational function f on X define its *order* $(a_1, \dots, a_n)_F \in \mathbb{Z}^n$ at the flag F . First let a_n be the order of f along X_{n-1} . We can write

$$f = f^{(n-1)} u_n^{a_n}, \quad a_n \in \mathbb{Z}.$$

Let $\tilde{f}^{(n-1)}$ be the restriction of $f^{(n-1)}$ to X_{n-1} , and a_{n-1} be the order of this restriction along X_{n-2} ,

$$\tilde{f}^{(n-1)} = \tilde{f}^{(n-2)} u_{n-1}^{a_{n-1}}, \quad a_{n-1} \in \mathbb{Z},$$

and so on. Finally,

$$\tilde{f}^{(1)} = f^{(0)} u_1^{a_1}, \quad a_1 \in \mathbb{Z}, \quad (\text{A2})$$

where a_1 is the order of $\tilde{f}^{(1)}$ at X_0 .

DEFINITION A.1. Let f_0, \dots, f_n be rational functions on X . Fix a complete flag F of irreducible subvarieties (A1). Let $(a_{i1}, \dots, a_{in})_F$ be the order of f_i at F . Denote $A = (a_{ij}) \in M_{n+1, n}(\mathbb{Z})$. Parshin's tame symbol of f_0, \dots, f_n at the flag F is the following non-zero element of k :

$$\langle f_0, \dots, f_n \rangle_F = (-1)^B \left(\prod_{i=0}^n f_i^{(-1)^i A_i} \right) (X_0), \quad (\text{A3})$$

where A_i is the determinant of the matrix obtained from A by eliminating its i th row, and

$$B = \sum_k \sum_{i < j} a_{ik} a_{jk} A_{ij}^k,$$

where A_{ij}^k is the determinant of the matrix obtained from A by eliminating its i th and j th rows and its k th column.

Note that the order of the rational function inside the large brackets in (A3) is $(0, \dots, 0)_F$, hence, its value at X_0 makes sense and is not zero.

Remark A.2. Let us associate with every rational function f on X a monomial $c u_1^{a_1} \dots u_n^{a_n}$, where $(u_1, \dots, u_n)_F$ are local parameters at F , $(a_1, \dots, a_n)_F$ is the order of f at F , and $c = f^{(0)}(X_0)$. Then the tame symbol $\langle f_0, \dots, f_n \rangle_F$ is equal to the symbol of the corresponding $n+1$ monomials (see Definition 3.8).

Parshin's symbol does not depend on the choice of local parameters $(u_1, \dots, u_n)_F$. It is multiplicative and skew-symmetric (compare to Proposition 3.9).

Example 1. Let f_1, \dots, f_n be rational functions on an algebraic variety X , whose zero loci $\{f_i = 0\}$ intersect transversely at a non-singular point $x \in X$. Denote by Z_i the irreducible component of $\{f_i = 0\}$ that contains x .

Let f_0 be any rational function on X whose divisor does not contain x . Then

$$\langle f_0, \dots, f_n \rangle_F = f_0(x)^{\mu(x)},$$

where $F : x = X_0 \subset X_1 \subset \dots \subset X_{n-1} \subset X$ for $X_i = Z_{i+1} \cap \dots \cap Z_n$, and $\mu(x) = \mu_1 \dots \mu_n$ is the product of the multiplicities μ_i of f_i along Z_i .

Indeed, for the system of local parameters at F we can choose the local equations of Z_i at x . Then the first row of the matrix A is zero and the other n rows form a lower triangular matrix with the multiplicities μ_i on the diagonal. Therefore, $A_0 = \mu_1 \dots \mu_n$ and $A_j = 0$, $1 \leq j \leq n$. It is also not hard to see that $B = 0$ (for example, one can use the description of B given in the proof of Proposition 3.9).

A.2 Parshin's residue

Let X be a complete algebraic variety over an algebraically closed field k , and F a complete flag of irreducible subvarieties (A1). Let $(u_1, \dots, u_n)_F$ be a system of local parameters at F , as before.

Consider a rational differential n -form on X . At a generic point of X_{n-1} the differentials du_1, \dots, du_n are linearly independent, and we can write

$$\omega = f du_1 \wedge \dots \wedge du_n, \quad \text{where} \quad f = \sum_{i_n > N_n} f_{i_n} u_n^{i_n}.$$

The restriction of the form $f_{-1} du_1 \wedge \dots \wedge u_{n-1}$ onto X_{n-1} makes sense and gives us a rational $(n-1)$ -form ω_{n-1} on X_{n-1} . Continuing in this way we arrive at a sequence of rational $(n-i)$ -forms ω_{n-i} on X_{n-i} , $i = 1, \dots, n$, the last one being a number $\omega_0 = f_{-1, \dots, -1}$ at the point X_0 . Note also

that this number is the coefficient of the series

$$f = \sum_{i_n \geq N_n} \sum_{i_{n-1} \geq N_{n-1}(i_n)} \cdots \sum_{i_1 \geq N_1(i_2, \dots, i_n)} f_{i_1, \dots, i_n} u_1^{i_1} \cdots u_n^{i_n}, \quad f_{i_1, \dots, i_n} \in k,$$

where we identify f with an element of the field $k((u_1)) \cdots ((u_n))$. Here $K((t))$ denotes the field of the Laurent power series in t with coefficients in a field K .

DEFINITION A.3. Let ω be a rational n -form on X . Fix a complete flag F of irreducible subvarieties (A1). Parshin's residue $\text{res}_F \omega$ at the flag F is the number $f_{-1, \dots, -1}$ constructed above.

Parshin's residue does not depend on the choice of local parameters $(u_1, \dots, u_n)_F$. The proof of this statement is similar to the proof we gave for the invariance of the toroidal residue in Proposition 4.2.

Example 2. Let f_1, \dots, f_n be rational functions on an algebraic variety X , whose zero loci $\{f_i = 0\}$ intersect transversely at a non-singular point $x \in X$. Denote by Z_i the irreducible component of $\{f_i = 0\}$ that contains x .

Let f_0 be any rational function on X which is regular in an open neighborhood of x . Then

$$\text{res}_F f_0 \frac{df_1}{f_1} \wedge \cdots \wedge \frac{df_n}{f_n} = \mu(x) f_0(x),$$

where $F : x = X_0 \subset X_1 \subset \cdots \subset X_{n-1} \subset X$, for $X_i = Z_{i+1} \cap \cdots \cap Z_n$, and $\mu(x) = \mu_1 \cdots \mu_n$ is the product of the multiplicities μ_i of f_i along Z_i .

Indeed, for the system of local parameters at F we can choose the local equations u_i of Z_i at x . Then

$$\text{res}_F f_0 \frac{df_1}{f_1} \wedge \cdots \wedge \frac{df_n}{f_n} = \text{res}_F f_0 \frac{du_1^{\mu_1}}{u_1^{\mu_1}} \wedge \cdots \wedge \frac{du_n^{\mu_n}}{u_n^{\mu_n}} = \text{res}_F \mu(x) f_0 \frac{du_1}{u_1} \wedge \cdots \wedge \frac{du_n}{u_n} = \mu(x) f_0(x).$$

A.3 Reciprocity laws

Now we will formulate Parshin's reciprocity laws for the tame symbol and the residue.

THEOREM A.4. Let X be a complete irreducible n -dimensional algebraic variety over an algebraically closed field k . Fix a partial flag of irreducible subvarieties $X_0 \subset \cdots \subset \widehat{X}_i \subset \cdots \subset X_n = X$, where X_i is omitted. Then

i) for any $n+1$ rational functions f_0, \dots, f_n on X

$$\prod_{X_i} \langle f_0, \dots, f_n \rangle_{X_0 \subset \cdots \subset X_i \subset \cdots \subset X_n} = 1,$$

ii) for any rational n -form ω on X

$$\sum_{X_i} \text{res}_{X_0 \subset \cdots \subset X_i \subset \cdots \subset X_n} \omega = 0,$$

where the product (sum) is taken over all irreducible subvarieties X_i that complete the flag, and is finite.

It follows by definition that Parshin's symbol $\langle f_0, \dots, f_n \rangle_F$ equals 1 unless F consists of intersections of components of the divisors of f_0, \dots, f_n . This shows that the above product is finite. Similarly, Parshin's residue $\text{res}_F \omega$ is zero unless F consists of intersections of components of the polar set of ω , hence, the above sum is finite.

In the proof of the main theorems (Theorems 3.15 and 4.8) we referred to the special case of the reciprocity law when $i = 0$. In this case the proof of the first part of Theorem A.4 is based on

the ‘reduction formula’ for the symbol. Namely, the property of the symbol analogous to the cofactor expansion for the determinant allows one to represent the n -dimensional symbol as a product of symbols of dimension $n - 1$ (see [FP04]). Then the statement follows from Weil’s reciprocity by induction.

For the case of the residue, notice that the residue of the n -form ω at the flag F is equal to the sum of the residues of 1-forms ω_1 on X_1 at X_0 . The statement then follows from the one-dimensional residue formula.

REFERENCES

- Ber75 D. N. Bernstein, *The number of roots of a system of equations*, Funct. Anal. Appl. **9** (1975), 183–185.
- BM96 J.-L. Brylinski and D. A. McLaughlin, *Multidimensional reciprocity laws*, J. reine angew. Math. **481** (1996), 125–147.
- BZ88 Y. D. Burago and V. A. Zalgaller, *Geometric inequalities*, Grundlehren Math. Wiss., vol. 285 (Springer, Berlin, 1988).
- CCD97 E. Cattani, D. Cox and A. Dickenstein, *Residues in toric varieties*, Compositio Math. **108** (1997), 35–76.
- CD97 E. Cattani and A. Dickenstein, *A global view of residues in the torus*, J. Pure Appl. Algebra **117/118** (1997), 119–144.
- CDS98 E. Cattani, A. Dickenstein and B. Sturmfels, *Residues and resultants*, J. Math. Sci. Univ. Tokyo **5** (1998), 119–148.
- Cox96 D. Cox, *Toric residues*, Ark. Mat. **34** (1996), 73–96.
- FP04 T. Fimmel and A. N. Parshin, *An introduction to the higher adelic theory*, book in preparation.
- Ful93 W. Fulton, *Introduction to toric varieties* (Princeton University Press, Princeton, NJ, 1993).
- Gel96 O. A. Gel’fond, *Combinatorial coefficients and the mixed volume of polytopes*, Funct. Anal. Appl. **30** (1996), 207–208.
- GK96 O. A. Gel’fond and A. G. Khovanskii, *Newton polyhedra and Grothendieck residues*, Dokl. Akad. Nauk **350** (1996), 298–300 (in Russian).
- GK02 O. A. Gel’fond and A. G. Khovanskii, *Toric geometry and Grothendieck residues*, Mosc. Math. J. **2** (2002), 99–112.
- Dan78 V. I. Danilov, *The geometry of toric varieties*, Russian Math. Surveys **33** (1978), 97–154.
- Kho99 A. G. Khovanskii, *Newton polyhedra, a new formula for mixed volume, product of roots of a system of equations*, Fields Inst. Commun. **24** (1999), 325–364.
- KKMS73 G. Kempf, F. Knudsen, D. Mumford and B. Saint-Donat, *Toroidal embeddings I*, Lecture Notes in Mathematics, vol. 339 (Springer, Berlin, 1973).
- Par75 A. N. Parshin, *Class fields and algebraic K-theory*, Uspekhi Mat. Nauk **30** (1975), 253–254.
- Par83 A. N. Parshin, *Chern classes, adeles and L-functions*, J. reine angew. Math., **341** (1983), 174–192.
- PS93 P. Pedersen and B. Sturmfels, *Product formulas for resultants and Chow forms*, Math. Z. **214** (1993), 377–396.
- Ser88 J.-P. Serre, *Algebraic groups and class fields*, Graduate Texts in Mathematics, vol. 117 (Springer, Berlin, 1988).
- Sop02 I. Soprounov, *Parshin’s symbols and residues, and Newton polyhedra*, PhD thesis, University of Toronto (2002).