



Cleveland State University  
**EngagedScholarship@CSU**

---

Mathematics Faculty Publications

Mathematics Department

---

1-1-1994

## Asymptotic Behavior of Solutions of Model Problems for a Coupled System

S. Shao

Cleveland State University, [s.shao@csuohio.edu](mailto:s.shao@csuohio.edu)

Follow this and additional works at: [https://engagedscholarship.csuohio.edu/scimath\\_facpub](https://engagedscholarship.csuohio.edu/scimath_facpub)

 Part of the [Mathematics Commons](#)

[How does access to this work benefit you? Let us know!](#)

---

### Repository Citation

Shao, S., "Asymptotic Behavior of Solutions of Model Problems for a Coupled System" (1994).

*Mathematics Faculty Publications*. 330.

[https://engagedscholarship.csuohio.edu/scimath\\_facpub/330](https://engagedscholarship.csuohio.edu/scimath_facpub/330)

This Article is brought to you for free and open access by the Mathematics Department at EngagedScholarship@CSU. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of EngagedScholarship@CSU. For more information, please contact [library.es@csuohio.edu](mailto:library.es@csuohio.edu).

# Asymptotic Behavior of Solutions of Model Problems for a Coupled System

S. SHAO

We study the asymptotic behavior of solutions of model problems for a coupled system. By consistent use of a priori estimates and asymptotic analysis, we present here an more efficient approach which provides precise descriptions of the asymptotic behavior of solutions of this system. Our results amplify and extend earlier results of Dorr, Parter, and Shampine and their treatment of this system. Meanwhile, the stability of the steady-state solutions of the corresponding time-dependent system is discussed. © 1994 Academic Press, Inc.

## 1. INTRODUCTION

Consider a Dirichlet problem for the coupled system of equations

$$\begin{aligned}u'' &= v && \text{in } (0, 1), \\u(0, \varepsilon) &= 0, && u(1, \varepsilon) = 0, \\ \varepsilon v'' + f(u, u')v' - g(x, u, u')v &= 0 && \text{in } (0, 1), \\v(0, \varepsilon) &= v_0, && v(1, \varepsilon) = v_1,\end{aligned}\tag{1.1}$$

for  $g(x, u, u') \geq 0$  and  $f(u, u') = h(u)$  or  $h(u')$ , where  $h(0) = 0$ . We assume that for each  $\varepsilon > 0$ , there exists at most one interior turning point  $x_0$  in  $(0, 1)$  such that  $u(x_0, \varepsilon) = 0$ . Throughout this paper we also assume that for  $0 \leq x \leq 1$ ,

- (i)  $f(u, u'')$  and  $g(x, u, u')$  are continuously differentiable functions of all variables and
- (ii)  $\partial f / \partial u$  and  $\partial f / \partial u'$ , do not change sign.

We are particularly interested in the case for which (i)  $dh/dz > 0$  or (ii)  $dh/dz < 0$  in  $(0, 1)$ .

System (1.1) is a simple model of the stream function-vorticity equations for two-dimensional, viscous, incompressible flow [8]. Historically this problem was considered by Dorr, Parter, and Shampine [3, 4]. They used analytical and numerical methods to study the existence of solutions of (1.1) and their asymptotic behavior as  $\varepsilon \rightarrow 0^+$  for the cases  $f(u, u') = u$  or  $u'$ . Howes and Shao [8] give an intuitive discussion of the system (1.1) for the case  $g \equiv 0$ . The purpose of this paper is to provide an alternative approach, which gives precise descriptions of the asymptotic behavior as  $\varepsilon \rightarrow 0^+$  of the solutions of system (1.1) for all boundary values of  $v$  and proves the existence of the limiting solutions of system (1.1) utilizing differential inequality techniques. Our results amplify and extend the earlier results of Dorr *et al.* and our treatment via asymptotic analysis is more appealing on both intuitive and analytical grounds. We hope that our results will be useful in the study of more complicated systems of ordinary and partial differential equations, such two-dimensional versions of system (1.1), etc.

The outline of this paper is as follows. In Section 2 we state some results from singular perturbation theory and some basic techniques of differential inequalities used later. The solutions of Model I,  $u'' = v$ ,  $\varepsilon v'' + h(u)v' = 0$ , Model II,  $u'' = v$ ,  $\varepsilon v'' + h(u')v' = 0$ , and Model III,  $u'' = v$ ,  $\varepsilon v'' + f(u, u')v' - g(x, u, u')v = 0$ , are discussed in Sections 3, 4, and 5, respectively. Finally, we discuss the stability of the steady-state solutions of the corresponding time-dependent system in Section 6.

## 2. PRELIMINARIES

We first state some elementary results of the theory of a singularly perturbed boundary value problem. Let us consider a linear model problem

$$\begin{aligned} \varepsilon y'' + f(x)y' &= 0, & x \text{ in } (0, 1), k \text{ is a constant,} & \quad (\text{LP}) \\ y(0, \varepsilon) &= A, & y(1, \varepsilon) &= B. \end{aligned}$$

Then as  $\varepsilon \rightarrow 0^+$ , we have the following classical results (cf. [5, Chap. 3; 1, Chap. 4]):

- (i)  $y(x, \varepsilon) \rightarrow B + BL(0)$ ,  $BL(0) = O(|A - B| \exp[-kx/\varepsilon])$  if  $f(x) \geq k > 0$  in  $[0, 1]$ ;
- (ii)  $y(x, \varepsilon) \rightarrow A + BL(1)$ ,  $BL(1) = O(|B - A| \exp[-k(1-x)/\varepsilon])$  if  $f(x) \leq -k < 0$  in  $[0, 1]$ ;
- (iii) if  $\exists x_0$  in  $(0, 1)$  such that  $f(x_0) = 0$  with  $f'(x_0) \neq 0$  (assume  $A < 0 < B$ ), then

- (a) If  $f'(x_0) < 0$ , then  $y(x, \varepsilon) \rightarrow B + BL(0)$  if  
 $x_0 > 1/2$ ,  $y(x, \varepsilon) \rightarrow A + BL(1)$  if  $x_0 < 1/2$ ;
- (b)  $y(x, \varepsilon) \rightarrow (A + B)/2 + BL(0) + BL(1)$   
if  $f'(x_0) < 0$  in  $[0, 1]$  and  $x_0 = 1/2$ ;
- (c)  $y(x, \varepsilon) \rightarrow \begin{cases} A \\ B \end{cases} + IL(x_0)$  if  $f'(x_0) > 0$  in  $[0, 1]$ .

We note that the asymptotic behavior of solutions of (LP) depends only on the signs of the coefficient  $f(x)$  of the first derivative  $y'$  and the boundary values  $A$  and  $B$ . This leads us to study the sign of the coefficient  $f(u, u')$  of the first derivative  $v'$  together with the boundary values of  $v$ .

A basic technique of differential inequalities is contained in the generalized Nagumo Theorem (cf. [4]). Applying the generalized Nagumo Theorem to our model problem (1.1), we only need to find bounding functions  $\alpha_i$  and  $\beta_i$  ( $i = 1, 2$ ) such that

$$\begin{aligned} \alpha_i &\leq \beta_i, & i = 1, 2, \\ \alpha_1(0, \varepsilon) &\leq 0 \leq \beta_1(0, \varepsilon), & \alpha_1(1, \varepsilon) \leq 0 \leq \beta_1(1, \varepsilon), \\ \alpha_1'' &\geq \beta_2, & \beta_1'' \leq \alpha_2, \\ \alpha_2(0, \varepsilon) &\leq v_0 \leq \beta_2(0, \varepsilon), & \alpha_2(1, \varepsilon) \leq v_1 \leq \beta_2(1, \varepsilon), \\ \varepsilon \alpha_2'' + f(u, z) \alpha_2' - g(x, u, z) \alpha_2 &\geq 0, \\ \varepsilon \beta_2'' + f(u, z) \beta_2' - g(x, u, z) \beta_2 &\leq 0, \end{aligned} \quad (2.1)$$

for all  $u$  in  $[\alpha_1, \beta_1]$  and  $z$  in  $\mathbb{R}$ . Then the system (1.1) has a unique solution  $u = u(x, \varepsilon)$ ,  $v = v(x, \varepsilon)$  such that

$$\alpha_1(x, \varepsilon) \leq u(x, \varepsilon) \leq \beta_1(x, \varepsilon), \quad \alpha_2(x, \varepsilon) \leq v(x, \varepsilon) \leq \beta_2(x, \varepsilon), \quad (2.2)$$

for  $x$  in  $[0, 1]$ .

**LEMMA.** *If we choose  $\alpha_1(x, \varepsilon) = Mx(x-1)/2$ ,  $\beta_1(x, \varepsilon) = mx(x-1)/2$ ,  $\alpha_2(x, \varepsilon) = m$ ,  $\beta_2(x, \varepsilon) = M$ , then the solution  $(u(x, \varepsilon), v(x, \varepsilon))$  of (1.1) satisfies the inequalities*

$$\frac{Mx(x-1)}{2} \leq u(x, \varepsilon) \leq \frac{mx(x-1)}{2}, \quad m \leq v(x, \varepsilon) \leq M, \quad (2.3)$$

for all  $x$  in  $[0, 1]$  and for all  $\varepsilon > 0$ , where

$$\begin{aligned} M &= \max\{v_0, v_1\}, \text{ and } m = \min\{v_0, v_1\} \text{ if } g(x, u, u') \equiv 0; \\ M &= \max\{0, v_0, v_1\}, \text{ and } m = \min\{0, v_0, v_1\} \text{ if } g(x, u, u') > 0. \end{aligned}$$

We divide system (1.1) into three models:

$$\text{Model I, } u'' = v, \varepsilon v'' + h(u)v' = 0,$$

$$\text{Model II, } u'' = v, \varepsilon v'' + h(u')v' = 0, \text{ and}$$

$$\text{Model III, } u'' = v, \varepsilon v'' + f(u, u')v' - g(x, u, u')v = 0.$$

Without lost generality, we assume  $dh/du > 0$  or  $dh/du' > 0$ .

### 3. MODEL I: $u'' = v, \varepsilon v'' + h(u)v = 0$

Consider the system

$$\begin{aligned} u'' &= v & \text{in } (0, 1), \\ u(0, \varepsilon) &= u(1, \varepsilon) = 0, \\ \varepsilon v'' + h(u)v' &= 0 & \text{in } (0, 1), \\ v(0, \varepsilon) &= v_0, & v(1, \varepsilon) = v_1. \end{aligned} \quad (3.1)$$

**THEOREM 1.** *Let  $(u(x, \varepsilon), v(x, \varepsilon))$  be the solution of (3.1). Then as  $\varepsilon \rightarrow 0^+$ , we have*

$$\frac{Mx(x-1)}{2} \leq u(x, \varepsilon) \leq \frac{mx(x-1)}{2} \quad \text{in } [0, 1];$$

$$(i) \quad v(x, \varepsilon) \rightarrow v_0 + BL(1) \text{ in } [0, 1] \text{ if } v_0 \geq 0, v_1 \geq 0;$$

$$(ii) \quad v(x, \varepsilon) \rightarrow v_1 + BL(0) \text{ in } [0, 1] \text{ if } v_0 \leq 0, v_1 \leq 0;$$

$$(iii) \quad v(x, \varepsilon) \rightarrow \begin{cases} v_0 \\ v_1 \end{cases} + IL(x_0) \text{ in } [0, 1] \text{ if } v_0 > 0, v_1 < 0; \text{ and}$$

$$(iv) \quad v(x, \varepsilon) \rightarrow 0 + BL(0) + BL(1) \text{ in } [0, 1] \text{ if } v_0 < 0, v_1 > 0,$$

where  $M = \max\{v_0, v_1\}$ ,  $m = \min\{v_0, v_1\}$ ,  $BL(k) =$  the boundary layer at  $x = k$ ,  $k = 0, 1$ .

*Proof.* We need to find bounding functions  $\alpha_i$  and  $\beta_i$  ( $i = 1, 2$ ) such that

$$\alpha_i \leq \beta_i, \quad i = 1, 2, \quad (E1)$$

$$\alpha_1(0, \varepsilon) \leq 0 \leq \beta_1(0, \varepsilon), \quad \alpha_1(1, \varepsilon) \leq 0 \leq \beta_1(1, \varepsilon), \quad (BC1)$$

$$\alpha_1'' \geq \beta_2, \quad \beta_1'' \leq \alpha_2, \quad (E2)$$

$$\varepsilon \alpha_2'' + h(u)\alpha_2' \geq 0, \quad \varepsilon \beta_2'' + h(u)\beta_2' \leq 0, \text{ for all } u \text{ in } [\alpha_1, \beta_1] \quad (E3)$$

$$\alpha_2(0, \varepsilon) \leq v_0 \leq \beta_2(0, \varepsilon), \quad \alpha_2(1, \varepsilon) \leq v_1 \leq \beta_2(1, \varepsilon). \quad (BC2)$$

Case 1.  $v_0 \geq 0, v_1 \geq 0$ . If  $v_0 = v_1 = 0$ , let  $\alpha_i = \beta_i = 0$  ( $i = 1, 2$ ), then it is clear that  $u(x, \varepsilon) \equiv 0$  and  $v(x, \varepsilon) \equiv 0$  is the solution. If  $v_0 = v_1 = p$ , let  $\alpha_2 = \beta_2 \equiv p$  and it follows that  $v(x, \varepsilon) \equiv p$ . For  $0 \neq v_0 \neq v_1 \neq 0$ , since  $u'' = v$ ,  $u(0, \varepsilon) = u(1, \varepsilon) = 0$ , we have  $u(x, \varepsilon) = v_0 x(x-1)/2$  and  $u < 0$  in  $(0, 1)$  and so  $h(u) < 0$  in  $(0, 1)$ . By virtue of the linear case, the solution  $v(x, \varepsilon)$  must display a boundary layer at  $x = 1$ .

We note that the reduced form of the  $v$ -equation  $h(u)v' = 0$  implies  $v = \text{constant}$  in  $(0, 1)$ . Therefore the constant is  $v_0$ . We set

$$\begin{aligned} \alpha_2(x, \varepsilon) &= v_0 - |v_1 - v_0| \exp\left[\frac{\lambda(1-x)}{\varepsilon}\right], \\ \beta_2(x, \varepsilon) &= v_0 + |v_1 - v_0| \exp\left[\frac{\lambda(1-x)}{\varepsilon}\right], \\ \alpha_1(x, \varepsilon) &= Mx(x-1), \quad \beta_1(x, \varepsilon) = \frac{mx(x-1)}{4}, \end{aligned} \quad (3.2)$$

where  $M = \max\{v_0, v_1\}$ ,  $m = \min\{v_0, v_1\}$ , and  $\lambda$  is a constant such that  $0 > \lambda > h(u)$  in  $[\delta, 1 - \delta]$  for a given  $\delta > 0$  and for  $\varepsilon > 0$ . Then the conditions (E1), (E2), (E3), (BC1), and (BC2) are satisfied. Therefore we have

$$\begin{aligned} &v_0 - |v_1 - v_0| \exp\left[\frac{\lambda(1-x)}{\varepsilon}\right] \\ &\leq v(x, \varepsilon) \leq v_0 + |v_1 - v_0| \exp\left[\frac{\lambda(1-x)}{\varepsilon}\right] \quad \text{in } [\delta, 1 - \delta], \end{aligned}$$

i.e.,  $v(x, \varepsilon) \rightarrow v_0 + BL(1)$  in  $[0, 1]$  as  $\varepsilon \rightarrow 0^+$ .

If we look at a neighborhood of  $x = 1$ , then

$$u(x, \varepsilon) \sim \frac{jx(x-1)}{2} \sim j(x-1), \quad (3.3)$$

where

$$j = \begin{cases} v_0 & \text{if } v_0 > 0, \\ v_1 & \text{if } v_0 = 0. \end{cases}$$

Substituting (3.3) into the equation  $\varepsilon v'' + h(u)v' = 0$ , we have

$$\varepsilon v'' + h\left(\frac{j(x-1)}{2}\right)v' = 0 \quad \text{in } (0, 1). \quad (3.4)$$

Introducing a stretched variable  $\xi = (1/\varepsilon)^{1/2} \left\{ \int h[j(x-1)/2] dx \right\}^{1/2}$  and setting  $w(\xi, \varepsilon) = v(x, \varepsilon)$ , then

$$w_{\xi\xi} + \xi w_{\xi} = 0, \quad w(0, \varepsilon) = v_1 \quad (3.5)$$

and we have

$$\begin{aligned} w(x, \varepsilon) &= v_1 \left[ 1 + \frac{2}{\pi^{1/2}} \int_0^{\xi/\sqrt{2}} \exp(-s^2) ds \right] \\ &= v_1 \left[ 1 + \operatorname{erf} \left( \frac{\xi}{\sqrt{2}} \right) \right], \end{aligned} \quad (3.6)$$

where  $\operatorname{erf}[y] = (2/\pi^{1/2}) \int_0^y \exp(-s^2) ds$  is the error function satisfying

$$\lim_{y \rightarrow -\infty} \operatorname{erf}[y] = -1.$$

Therefore, the thickness of the boundary layer at  $x = 1$  is of order  $\varepsilon^{1/2}$ .

*Case 2.*  $v_0 \leq 0, v_1 \leq 0$ . In this case, by the same arguments as in Case 1, we have  $h(u) > 0$  in  $(0, 1)$ , and so the solution  $v(x, \varepsilon)$  of the  $v$ -equation will display a boundary layer at  $x = 0$ . We make the change of variables

$$y = 1 - x, \quad m(y, \varepsilon) = -u(1 - y, \varepsilon), \quad n(y, \varepsilon) = -v(1 - y, \varepsilon).$$

Then the system (3.1) becomes

$$\begin{aligned} m'' &= n, & m(0, \varepsilon) &= m(1, \varepsilon) = 0, \\ \varepsilon n'' + h(m)n' &= 0, & n(0, \varepsilon) &= n_0 = -v_1, & n(1, \varepsilon) &= n_1 = -v_0, \end{aligned}$$

provided  $n_0 \geq 0$  and  $m_1 \geq 0$ . By our previous results, we obtain

$$n(y, \varepsilon) \rightarrow n_0 + BL(1) \quad \text{in } [0, 1] \text{ as } \varepsilon \rightarrow 0^+, \text{ i.e., } v(x, \varepsilon) \rightarrow v_1 + BL(0).$$

The thickness of the boundary layer at  $x = 0$  is also of order  $\varepsilon^{1/2}$ .

*Case 3.*  $v_0 > 0, v_1 < 0$ . We note that  $u(x, \varepsilon)$  changes sign in  $(0, 1)$  in this case, and so does  $h(u)$ . Since  $u'' = v, u(0, \varepsilon) = u(1, \varepsilon) = 0$ , there exists a unique  $x_0$  in  $(0, 1)$  such that  $u(x_0, \varepsilon) = 0$  and hence  $h(u(x_0, \varepsilon)) = 0$ . Since  $u < 0$  and  $u > 0$  near  $x = 1$ , we have  $h(u) < 0$  near  $x = 0$  and  $h(u) > 0$  near  $x = 1$ . We claim that the solution  $v = v(x, \varepsilon)$  of (3.1) displays an interior layer as  $\varepsilon \rightarrow 0^+$  in  $(0, 1)$ , that is,

$$\lim_{\varepsilon \rightarrow 0^+} v(x, \varepsilon) = V(x) = \begin{cases} v_0 & \text{if } 0 \leq x \leq x_0 - \delta, \\ v_1 & \text{if } x_0 + \delta \leq x < 1 - \delta, \end{cases} \quad (3.7)$$

for  $x_0$  in  $(0, 1)$ ,  $0 < \delta < 1$ . Since  $\varepsilon v'' + h(u)v' = 0$  in  $(0, 1)$ , we have

$$\varepsilon \ln \frac{|v'(1, \varepsilon)|}{|v'(0, \varepsilon)|} = - \int_0^1 h(u(s, \varepsilon)) ds \quad \text{in } (0, 1). \quad (3.8)$$

We know that  $|v'(0, \varepsilon)| = |v'(1, \varepsilon)| = O(\varepsilon^{1/2})$  as  $\varepsilon \rightarrow 0^+$ ; it follows that

$$\lim_{\varepsilon \rightarrow 0^+} \varepsilon \ln \frac{|v'(1, \varepsilon)|}{|v'(0, \varepsilon)|} = 0 \quad \text{in } (0, 1).$$

Hence, we have

$$\int_0^1 h(u(s, \varepsilon)) ds \rightarrow 0 \quad \text{in } (0, 1) \text{ as } \varepsilon \rightarrow 0^+. \quad (3.9)$$

Let  $U(x) = \lim_{\varepsilon \rightarrow 0^+} u(x, \varepsilon)$ , then  $U'' = V$ ,  $U(0) = U(1) = 0$ . By (3.4), we obtain

$$U(x) = \begin{cases} v_0 x(x - x_0)/2, & \text{if } 0 \leq x \leq x_0, \\ -v_1 (1 - x)(x - x_0)/2 & \text{if } x_0 \leq x \leq 1 \end{cases} \quad (3.10)$$

and so there exists a constant  $k$  such that  $U(x) \geq k > 0$  for  $0 < x < x_0 - \delta$  and  $U(x) \leq -k < 0$  for  $x_0 + \delta < x < 1$ . It follows that  $h(U(x)) \geq r > 0$  in  $(0, x_0 - \delta)$  and  $h(U(x)) \leq -r < 0$  in  $(x_0 + \delta, 1)$ , where  $r$  is a positive constant. Therefore, there is an interior layer at  $x = x_0$  in  $(0, 1)$ . By the Dominated Convergence Theorem [9] and (3.9), it follows that as  $\varepsilon \rightarrow 0^+$ ,  $0 = \int_0^1 [h(u(s, \varepsilon)) \rightarrow 0] ds$ , which implies that

$$\begin{aligned} 0 &= \int_0^1 h(U(s)) ds \\ &= \int_0^{x_0} h(U(s)) ds + \int_{x_0}^1 h(U(s)) ds \quad \text{in } (0, 1), \end{aligned} \quad (3.11)$$

and also

$$\begin{aligned} v(x, \varepsilon) &= V + |v_1 - v_0| \left\{ 1 + \operatorname{erf} \left[ \frac{1}{\sqrt{2\varepsilon}} \int h \left( \frac{j}{2(x - x_0)} \right) dx \right] \right\} \\ &\quad \text{for } 0 \leq x \leq 1, \\ v(x, \varepsilon) &= V + IL(x_0), \quad 0 \leq x \leq 1, \end{aligned}$$

where  $v(x, \varepsilon)$  is the solution of (3.1),  $V = V(x)$  as in (3.7),  $IL(x_0)$  means the interior layer at  $x_0$  which is determined by (3.6).

*Case 4.*  $v_0 < 0, v_1 > 0$ . In this case  $u(x, \varepsilon)$  also changes sign in  $(0, 1)$  and there exists a unique  $x_0$  in  $(0, 1)$  such that  $h(u(x_0)) = 0$ . However, there is no interior layer in  $(0, 1)$  because of  $u(x, \varepsilon) > 0$  near  $x = 0$  and  $u(x, \varepsilon) < 0$  near  $x = 1$ . The wrong sign of  $u$  does not allow an interior layer to be displayed in  $(0, 1)$ . We consider the asymptotic behavior as  $\varepsilon \rightarrow 0^+$  of



solutions of (3.1) in the subintervals  $[0, x_0]$  and  $[x_0, 1]$  separately. Let  $v^* = v(x_0, \varepsilon)$ , then we have the systems of equations

$$\begin{aligned} u'' &= v && \text{in } (0, x_0), \\ u(0, \varepsilon) &= u(x_0, \varepsilon) = 0, \\ \varepsilon v'' + h(u)v' &= 0 && \text{in } (0, x_0), \\ v(0, \varepsilon) &= v_0, v(x_0, \varepsilon) = v^* \end{aligned} \tag{SI}$$

and

$$\begin{aligned} u'' &= v && \text{in } (x_0, 1), \\ u(x_0, \varepsilon) &= u(1, \varepsilon) = 0, \\ \varepsilon v'' + h(u)v' &= 0 && \text{in } (x_0, 1), \\ v(x_0, \varepsilon) &= v^*, \quad v(1, \varepsilon) = v_1. \end{aligned} \tag{SII}$$

For  $1 \gg \delta > 0$ , we have  $h(u) > 0$  in  $[\delta, x_0 - \delta]$  and  $h(u) < 0$  in  $[x_0 + \delta, 1]$ . Since  $v_0 < 0$ ,  $v^* = 0$ ,  $v_1 > 0$ , it follows that (SI) and (SII) are Case 2 and Case 1, respectively. Hence, by our previous results, we have that as  $\varepsilon \rightarrow 0^+$ ,

$$v(x, \varepsilon) \rightarrow v^* + BL(0) \text{ in } [\delta, x_0 - \delta]$$

and

$$v(x, \varepsilon) \rightarrow v^* + BL(1) \text{ in } [x_0 + \delta, 1 - \delta].$$

Since  $v^* = v(x_0, \varepsilon) = 0$ , it follows that  $v(x, \varepsilon) \rightarrow 0 + BL(0) + BL(1)$  in  $[0, 1]$  as  $\varepsilon \rightarrow 0^+$ .

This finishes the proofs for all choices of the boundary data  $v_0$  and  $v_1$ . All of the previous results are summarized in Fig. 1.




FIG. 1. Boundary value portrait for model I.

#### 4. THE SOLUTIONS OF MODEL II: $u'' = v, \varepsilon v'' + h(u')v = 0$

We consider the problem

$$\begin{aligned} u'' &= v && \text{in } (0, 1), \\ u(0, \varepsilon) &= u(1, \varepsilon) = 0, \\ \varepsilon v'' + h(u')v' &= 0 && \text{in } (0, 1), \\ v(0, \varepsilon) &= v_0, && v(1, \varepsilon) = v_1. \end{aligned} \tag{4.1}$$

There is a fundamental difference between the systems (4.1) and (3.1). Because  $u'(x, \varepsilon)$  always changes sign in  $(0, 1)$ , even if  $u(x, \varepsilon)$  does not change sign, then there is at least *one* interior turning point  $x_0$  in  $(0, 1)$  such that  $h(u'(x_0, \varepsilon)) = 0$ . The behavior of the coefficient  $h(u')$  of  $v'$  in the  $v$ -equation of (4.1) is more difficult to determine than the behavior of  $h(u)$  in (3.1). We give an intuitive analysis of the asymptotic behavior as  $\varepsilon \rightarrow 0^+$  of the solution of (4.1). If  $v_0 = v_1$ , then  $v(x, \varepsilon) \equiv v_0$  in  $[0, 1]$  and  $u(x, \varepsilon) = v_0 x(x-1)/2$  in  $[0, 1]$  [cf. Section 3]. For the other boundary values  $v_0$  and  $v_1$ , we examine the following four cases.

*Case 1.*  $v_0 \geq 0, v_1 \geq 0$ . If  $v_0 > 0, v_1 > 0$ , then there exists some  $x_0$  in  $(0, 1)$  such that  $u'(x_0) = 0$  and  $u'(x, \varepsilon) < 0$  near  $x = 0$ , while  $u'(x, \varepsilon) > 0$  near  $x = 1$ . It follows that  $h(u'(x_0)) = 0, h(u') < 0$  near  $x = 0$  and  $h(u') > 0$  near  $x = 1$ . Consequently, the sign of  $h(u')$  at each endpoint does not allow  $v(x, \varepsilon)$  to have boundary layers. The only possible asymptotic behavior available to  $v$  is then interior layer behavior at the point  $x_0$  in  $(0, 1)$ . Such behavior is indeed possible. If  $v_1 > v_0$  then  $v' > 0$  in  $(0, 1)$  and  $h(u') > 0$  near  $x = 1$ ; consequently,  $v'' < 0$  in  $(x_0, 1)$  by (4.3). Also we have  $v'' > 0$  in  $(0, x_0)$  since  $h(u') < 0$  near  $x = 0$  and so  $v(x, \varepsilon)$  must have an interior layer in  $(0, 1)$ . If  $v_0 > v_1$  then  $v' < 0$  in  $(0, 1)$ ,  $h(u') < 0$  near  $x = 0$ , and  $h(u') > 0$  near  $x = 1$ ; consequently,  $v'' > 0$  in  $(0, x_0)$  and  $v'' < 0$  in  $(x_0, 1)$ . This also implies that  $v(x, \varepsilon)$  has an interior layer in  $(0, 1)$ . To determine the location  $x_0$  of the interior layer, we let

$$U(x) = \lim_{\varepsilon \rightarrow 0^+} u(x, \varepsilon) \quad \text{for } x \neq x_0.$$

Then  $U(x)$  must satisfy the following (five) conditions:  $U(0) = U(1) = 0$ ,  $U(x_0^+) = U(x_0^-)$ , and  $U'(x_0^+) = U'(x_0^-) = 0$ . We know that  $U'' = v_0$  for  $x \leq x_0$  and  $U'' = v_1$  for  $x \geq x_0$ ; therefore,  $U(x) = v_0 x^2/2 + ax + b$  for  $x \leq x_0$ ,  $U(x) = v_1 x^2/2 + cx + d$  for  $x \geq x_0$ , where  $a, b, c$ , and  $d$  are constants. Since  $U(0) = U(1) = 0$ , we have that  $b = 0$  and  $c + d = -v_1/2$ , and the three remaining conditions allow us to determine  $a, c$ , and  $x_0$ . Since  $U'(x_0^+) = U'(x_0^-) = 0$  we find that  $a = -v_0 x_0$  and  $c = -v_1 x_0$ , that is,  $U(x) =$

$v_0 x(x - 2x_0)/2$  for  $x \leq x_0$ ,  $U(x) = v_1(x - 1)(x + 1 - 2x_0)/2$  for  $x \geq x_0$ . Finally, in order that  $U(x_0^+) = U(x_0^-)$ , we must have  $-v_0 x_0^2/2 = -v_1(1 - x_0)^2/2$ , that is,  $x_0 = v_1^{1/2}/(v_0^{1/2} + v_1^{1/2})$ .

*Case 2.*  $v_0 \leq 0, v_1 \leq 0$ . The reader will see immediately that this case is not the reflection of Case 1, so there is no change of variable of the type employed in showing the equivalence of Cases 1 and 2 for Model I. Since  $u'(x, \varepsilon) > 0$  near  $x = 0$ ,  $u'(x_0, \varepsilon) = 0$  for some point  $x_0$  in  $(0, 1)$  and  $u'(x, \varepsilon) < 0$  near  $x = 1$ . The coefficient  $h(u')$  of  $v'$  in the  $v$ -equation behaves accordingly. Consequently, the solution  $v(x, \varepsilon)$  in the  $v$ -equation cannot have an interior layer at  $x_0$ . However, the sign of  $h(u')$  at the endpoints is compatible with the existence of boundary layers in  $v(x, \varepsilon)$ . For if  $v_0 > v_1$  then  $v' < 0$ , since  $h(u') > 0$  near  $x = 0$  and  $h(u') < 0$  near  $x = 1$ , with of course  $v'' > 0$  near  $x = 0$  and  $v'' < 0$  near  $x = 1$  by (4.3). Similarly, if  $v_0 < v_1$  then  $v' > 0$ , and we must have  $v'' < 0$  near  $x = 0$  and  $v'' > 0$  near  $x = 1$  and near  $x = 0$ ,  $h(u') < 0$  near  $x = 1$ . Therefore,  $v(x, \varepsilon) \rightarrow c + BL(0) + BL(1)$  as  $\varepsilon \rightarrow 0^+$  with  $c$  a constant strictly between  $v_0$  and  $v_1$ . In order to determine  $c$  and the solution  $v(x, \varepsilon)$  of (4.1), we let  $U(x) = \lim_{\varepsilon \rightarrow 0^+} u(x, \varepsilon)$  in  $[0, 1]$  and  $c^* = \lim_{\varepsilon \rightarrow 0^+} \varepsilon[v'(1, \varepsilon) - v'(0, \varepsilon)]$ . Then we know that  $U'' = c$ , so  $U(x) = cx(x - 1)/2$ , giving us  $U'(x) = c(2x - 1)/2$  in  $(0, 1)$ ,  $U'(0) = -c/2$ ,  $U'(1) = c/2$ . It follows that  $U'(0) > 0$  and  $U'(1) < 0$  provided  $c \leq 0$ . We also have  $v(x, \varepsilon) \rightarrow c + (v_0 - c) \exp[-kx/\varepsilon] + (v_1 - c) \exp[-k(1 - x)/\varepsilon]$  as  $\varepsilon \rightarrow 0^+$  in  $[0, 1]$ , where  $k = |c|/2$ . From (4.3),  $\varepsilon v'' = -v'h(u')$ , we have  $-\varepsilon \int_0^1 v'' dx = \int_0^1 v'h(u') dx = v h(u')|_0^1 - \int_0^1 v dh(u')$ , which implies that

$$-\varepsilon[v'(1, \varepsilon) - v'(0, \varepsilon)] \rightarrow v_1 h(U'(1)) - v_0 h(U'(0)) - c \int_0^1 \left( \frac{\partial h}{\partial x} + U'' \frac{\partial h}{\partial u'} \right) dx \quad \text{as } \varepsilon \rightarrow 0^+,$$

that is,

$$c^* = c \left[ \frac{1}{2} (v_0 + v_1) + \int_0^1 \left( \frac{\partial h}{\partial x} + c \frac{\partial h}{\partial u'} \right) dx \right]; \quad (4.2)$$

hence,  $c$  must satisfy (4.2). If  $\int_0^1 h(u') dx = 0$ , then  $v'(0, \varepsilon) = v'(1, \varepsilon)$ , we have

$$\frac{-k(v_0 - c)}{\varepsilon} = \frac{k(v_1 - c)}{\varepsilon} \quad \text{or} \quad c = \frac{(v_0 + v_1)}{2}.$$

*Case 3.*  $v_0 < 0, v_1 > 0$ . We note that  $u' > 0$  near both endpoints and  $u'(x_0, \varepsilon) = 0$  for some  $x_0$  in  $(0, 1)$ , which implies that  $h(u') > 0$  near both endpoints. Hence,  $v(x, \varepsilon)$  has no interior layer in  $(0, 1)$ , and the sign of  $h(u')$  near  $x = 0$  allows  $v(x, \varepsilon)$  to have a boundary layer there. Meanwhile, since  $v' > 0$  (from the fact that  $v_1 > v_0$ ) in  $(0, 1)$ , we must have  $v'' < 0$  near




FIG. 2. Behavior of  $u$ .

$x = 1$  since  $h(u') > 0$  there, and so  $v(x, \epsilon)$  has an "S-layer" (interior layer at  $x = 1$ ) near  $x = 1$ . Therefore,  $v(x, \epsilon) \rightarrow c$  in  $[\delta, 1 - \delta^*]$  as  $\epsilon \rightarrow 0^+$ , where  $c$  is a constant determined by (4.2), with  $\delta > 0$  and  $\delta^* > 0$ . If  $\int_0^1 h(u') dx = 0$ , then  $\epsilon \ln(|v'(1, \epsilon)|/|v'(0, \epsilon)|) = 0$  or  $|v'(1, \epsilon)| = |v'(0, \epsilon)|$ , and so the only possible asymptotic behavior available to  $v(x, \epsilon)$  is twin layer behavior and  $c = (v_0 + v_1)/2$ ,  $c \leq 0$ . So if  $v_0 + v_1 > 0$ , then  $c$  must be equal to zero. The condition  $v_0 + v_1 > 0$  says that  $v_1 > |v_0|$ , and so there exists a unique point  $\xi(\epsilon)$  in  $(0, 1)$  such that  $u''(x, \epsilon) < 0$  for  $0 \leq x < \xi(\epsilon)$ ,  $u''(x, \epsilon) = 0$ , and  $u''(x, \epsilon) > 0$  for  $\xi(\epsilon) < x < 1$ , with  $\xi(\epsilon) \rightarrow 1$  as  $\epsilon \rightarrow 0^+$ ; cf. Fig. 2. (This point  $\xi$  is simply the unique point in  $(0, 1)$  where  $v(\xi(\epsilon), \epsilon) = 0$ ; its existence follows from the fact that  $v'(x, \epsilon) > 0$  in  $[0, 1]$ ). Therefore, in the boundary layer at  $x = 0$ ,  $v'' < 0$ , with  $h(u') > 0$ ; however, the boundary layer at  $x = 1$  is more complicated, in that  $\xi(\epsilon)$  is an inflection point for  $v$ . Consider  $\epsilon v'' = -h(u')v'$ , slightly to the left of  $\xi$ ,  $v'' > 0$  and  $h(u') < 0$ ; at  $x = \xi$ ,  $v''(\xi, \epsilon) = 0$  and  $h(u'(\xi, \epsilon)) = 0$ ; and to the right of  $\xi$ ,  $v'' < 0$  and  $h(u') > 0$ ; cf. Fig. 3.

Suppose next that  $v_0 + v_1 < 0$ . We will show that  $c = 0$  is impossible. If  $v \rightarrow 0$  in  $(0, 1)$  as  $\epsilon \rightarrow 0^+$ , then there exists a unique point  $\eta = \eta(\epsilon)$  in  $(0, 1)$  such that  $u''(x, \epsilon) < 0$  in  $[0, \eta(\epsilon))$ ,  $(u''(\eta, \epsilon)) = 0$  and  $u'' > 0$  in  $(\eta(\epsilon), 1]$  with  $\eta(\epsilon) \rightarrow 0$  as  $\epsilon \rightarrow 0^+$ ; cf. Fig. 4 (the existence of  $\eta$  follows from the same argument used in proving the existence of  $\xi$ ). In the boundary layer at  $x = 0$ ,  $v'' > 0$  in  $[0, \eta(\epsilon))$ ,  $v''(\eta, \epsilon) = 0$ , and  $v'' < 0$  for  $x$  slightly to the right of  $\eta$ . In




FIG. 3.  $v_0 + v_1 > 0$ .




FIG. 4. Behavior of  $u$ .

turn, we see that  $h(u') > 0$  in  $[0, \eta(\epsilon))$ ,  $h(u'(\eta, \epsilon)) = 0$  and  $h(u') < 0$  for  $x$  slightly to the right of  $\eta$ . These relations are *incompatible* with the relation  $\epsilon v'' = -h(u')v'$ . We conclude that for  $v_0 + v_1 < 0$ , the limiting value  $c$  of  $v(x, \epsilon)$  must be negative, that is,  $c = (v_0 + v_1)/2$ ; cf. Fig. 5. Actually, the exact solution of the  $v$ -equation  $\epsilon v'' + u'v' = 0$ , with  $v(0, \epsilon) = v_0$ ,  $v(1, \epsilon) = v_1$ , is

$$v(x, \epsilon) = v_0 + (v_1 - v_0) \frac{\left(\int_0^x \exp[-u(s, \epsilon)/\epsilon] ds\right)}{\left(\int_0^1 \exp[-u(s, \epsilon)/\epsilon] ds\right)}.$$

We can use Laplace's Method to obtain

$$\frac{\left\{\int_0^x \exp[-u(s, \epsilon)/\epsilon] ds\right\}}{\left\{\int_0^1 \exp[-u(s, \epsilon)/\epsilon] ds\right\}} \rightarrow \frac{1}{2} \quad \text{as } \epsilon \rightarrow 0^+$$

in  $(0, 1)$ , which implies that  $\lim_{\epsilon \rightarrow 0^+} v(x, \epsilon) = (v_0 + v_1)/2 + BL(0) + SL(1)$  for  $x$  in  $[0, 1]$ .

*Case 4.*  $v_0 > 0$ ,  $v_1 < 0$ . Fortunately the asymptotic behavior of  $v(x, \epsilon)$  in this case is a reflection of that observed in the previous case. So there is a "Z-layer" (backwards S-layer) displayed at  $x = 0$ . Thus  $v(x, \epsilon) \rightarrow c + BL(1) + ZL(0)$  as  $\epsilon \rightarrow 0^+$  in  $[0, 1]$ , where  $c$  is determined by (4.2). We




FIG. 5.  $v_0 + v_1 < 0$ .




FIG. 6. Boundary value portrait for model II.

summarize our results for (4.1) in the form of a boundary value portrait in Fig. 6.

5. SOLUTION OF MODEL III:  $u'' = v, \varepsilon v'' + f(u, u')v' - g(x, u, u')v = 0$

The problem we consider in this section is the system (1.1) in which  $g(x, u, u') \geq g^* > 0$ , for  $g^*$  a constant, that is,

$$\begin{aligned}
 u'' &= v & \text{in } (0, 1), \\
 u(0, \varepsilon) &= u(1, \varepsilon) = 0, \\
 \varepsilon v'' + f(u, u')v' - g(x, u, u')v &= 0 & \text{in } (0, 1), \\
 v(0, \varepsilon) &= v_0, & v(1, \varepsilon) = v_1,
 \end{aligned} \tag{5.1}$$

where  $f(u, u') = h(u)$  or  $h(u')$  and  $g(x, u, u') \in C^1[0, 1] \times \mathbb{R}^2$ . The strict positivity of the function  $g(x, u, u')$  in the  $v$ -equation forces the asymptotic behavior as  $\varepsilon \rightarrow 0^+$  of the solution  $v(x, \varepsilon)$  to be independent of the coefficient  $f(u, u')$  of  $v'$ , because the reduced form of the  $v$ -equation,  $f(u, u')v' + g(x, u, u')v = 0$ , has only the zero solution  $v \equiv 0$  in  $(0, 1)$ ; cf. [3]. We have the following theorem.

**THEOREM 2.** *Let  $v(x, \varepsilon)$  be the solution of (6.1), then as  $\varepsilon \rightarrow 0^+$ ,*

$$\begin{aligned}
 v(x, \varepsilon) &\rightarrow 0 + BL(0) + BL(1) \\
 &\text{in } [0, 1] \text{ for all boundary data } v_0 \neq 0 \text{ and } v_1 \neq 0, \\
 v(x, \varepsilon) &\rightarrow 0 + BL(1) \quad \text{for } v_0 \neq 0, v_1 > 0
 \end{aligned}$$

and

$$v(x, \varepsilon) \rightarrow 0 + BL(0) \quad \text{for } v_0 > 0, v_1 \neq 0.$$

*Proof.* The basic technique we use is the construction of bounding functions  $\alpha_i$  and  $\beta_i$  ( $i = 1, 2$ ) such that

$$\alpha_i \leq \beta_i, \quad i = 1, 2, \quad (\text{A1})$$

$$\alpha_1(0, \varepsilon) \leq 0 \leq \beta_1(0, \varepsilon), \quad \alpha_1(1, \varepsilon) \leq 0 \leq \beta_1(1, \varepsilon), \quad (\text{AB1})$$

$$\alpha_1'' \geq \beta_2, \quad \beta_1'' \leq \alpha_2, \quad (\text{A2})$$

$$\alpha_2(0, \varepsilon) \leq v_0 \leq \beta_2(0, \varepsilon), \quad \alpha_2(1, \varepsilon) \leq v_1 \leq \beta_2(1, \varepsilon), \quad (\text{AB2})$$

$$\begin{aligned} \varepsilon \alpha_2'' + f(u, z) \alpha_2' - g(x, u, z) \alpha_2 &\geq 0, \\ \varepsilon \beta_2'' + f(u, z) \beta_2' - g(x, u, z) \beta_2 &\leq 0, \end{aligned} \quad (\text{A3})$$

for all  $u$  in  $[\alpha_1, \beta_1]$  and  $z \in \mathbb{R}$ . Again, we divide our discussion into four cases for the different signs of the boundary values  $v_0, v_1$  when  $f(u, u') = h(u)$  or  $h(u')$ . We note that  $v_0 = v_1 = 0$  implies  $v = 0$  and  $u = 0$ . We now consider the cases in which  $v_0$  and  $v_1$  are not both zero, and without loss of generality, we assume  $\partial h / \partial z \geq 0$  for  $z = u$  or  $u'$ .

*Part (I).*  $f(u, u') = h(u)$ . *Case 1.*  $v_0 \geq 0, v_1 \geq 0$ . We note that it is sufficient to show that  $\varepsilon \beta_2'' + h(\beta_1) \beta_2' - g^* \beta_2 \leq 0$  if  $\alpha_2(x, \varepsilon) = 0$ . We define, for  $0 \leq x \leq 1$  and  $\varepsilon > 0$ ,

$$\alpha_1(x, \varepsilon) = \frac{1}{2}(v_1 + v_0)(x^2 - x), \quad \beta_1(x, \varepsilon) = 0, \quad \alpha_2(x, \varepsilon) = 0,$$

and

$$\beta_2(x, \varepsilon) = v_0 \exp \left[ -\left(\frac{g^*}{\varepsilon}\right)^{1/2} x \right] + v_1 \exp \left[ -\left(\frac{g^*}{\varepsilon}\right)^{1/2} (1-x) \right].$$

Then it follows that (A1), (A2), and (AB1) are satisfied. Since  $h(0) = 0$ , that is,  $h(\beta_1) = 0$ , it follows that

$$\begin{aligned} &\varepsilon \beta_2'' + h(\beta_1) \beta_2' - g^* \beta_2 \\ &= \varepsilon \left(\frac{g^*}{\varepsilon}\right) v_0 \exp \left[ -\left(\frac{g^*}{\varepsilon}\right)^{1/2} x \right] \\ &\quad + \varepsilon \left(\frac{g^*}{\varepsilon}\right) v_1 \exp \left[ -\left(\frac{g^*}{\varepsilon}\right)^{1/2} (1-x) \right] \\ &\quad - g^* v_0 \exp \left[ -\left(\frac{g^*}{\varepsilon}\right)^{1/2} x \right] - g^* v_1 \exp \left[ -\left(\frac{g^*}{\varepsilon}\right)^{1/2} (1-x) \right] = 0 \end{aligned}$$

and

$$\alpha_1''(x, \varepsilon) = (v_1 + v_0) \geq \beta_2, \quad \beta_1'' = 0 = \alpha_2.$$

Therefore, we see that

$$\begin{aligned} 0 \leq v(x, \varepsilon) \leq v_0 \exp \left[ - \left( \frac{g^*}{\varepsilon} \right)^{1/2} x \right] \\ + v_1 \exp \left[ - \left( \frac{g^*}{\varepsilon} \right)^{1/2} (1-x) \right] \quad \text{in } [0, 1]. \end{aligned}$$

As  $\varepsilon \rightarrow 0^+$ , we have

$$\begin{aligned} v(x, \varepsilon) &\rightarrow 0 + BL(0) + BL(1) & \text{for } v_0 > 0, v_1 > 0, \\ v(x, \varepsilon) &\rightarrow 0 + BL(1) & \text{for } v_0 = 0, v_1 > 0, \end{aligned}$$

and

$$v(x, \varepsilon) \rightarrow 0 + BL(0) \quad \text{for } v_0 > 0, v_1 = 0.$$

*Case 2.*  $v_0 \leq 0, v_1 \leq 0$ . In this case, for  $x$  in  $[0, 1]$  and  $\varepsilon > 0$ , we define

$$\begin{aligned} \alpha_1(x, \varepsilon) &= 0, \quad \beta_1(x, \varepsilon) = \frac{1}{2}(v_1 + v_0)(x^2 - x), \\ \alpha_2(x, \varepsilon) &= -|v_0| \exp \left[ - \left( \frac{g^*}{\varepsilon} \right)^{1/2} x \right] \\ &\quad - |v_1| \exp \left[ - \left( \frac{g^*}{\varepsilon} \right)^{1/2} (1-x) \right] - \Gamma(\varepsilon), \quad \beta_2(x, \varepsilon) = 0, \end{aligned}$$

where  $\Gamma(\varepsilon) = \varepsilon\rho/g^*$ ,  $\rho$  is a positive constant. By the same calculations as in Case 1, we have

$$-|v_0| \exp \left[ - \left( \frac{g^*}{\varepsilon} \right)^{1/2} x \right] - |v_1| \exp \left[ - \left( \frac{g^*}{\varepsilon} \right)^{1/2} (1-x) \right] - \Gamma(\varepsilon) \leq v(x, \varepsilon) \leq 0,$$

from which it follows that

$$\begin{aligned} v(x, \varepsilon) &\rightarrow 0 + BL(0) + BL(1) & \text{for } v_0 < 0, v_1 < 0, \\ v(x, \varepsilon) &\rightarrow 0 + BL(1) & \text{for } v_0 = 0, v_1 < 0, \end{aligned}$$

and

$$v(x, \varepsilon) \rightarrow 0 + BL(0) \quad \text{for } v_0 < 0, v_1 = 0 \text{ as } \varepsilon \rightarrow 0^+ \text{ in } [0, 1].$$



Case 3.  $v_0 < 0$ ,  $v_1 > 0$ . Since  $u(x, \varepsilon)$  changes sign in this case, there exists a unique interior turning point  $x_0$  in  $(0, 1)$  such that  $u(x_0, \varepsilon) = 0$ , which implies that  $h(u(x_0)) = 0$  and  $v(x_0, \varepsilon) = 0$ . We consider the systems

$$\begin{aligned} u'' &= v && \text{in } (0, x_0), \\ u(0, \varepsilon) &= u(x_0, \varepsilon) = 0, \\ \varepsilon v'' + h(u)v' - g(x, u, u')v &= 0 && \text{in } (0, x_0), \\ v(0, \varepsilon) &= v_0, && v(x_0, \varepsilon) = v^* \end{aligned} \tag{T1}$$

and

$$\begin{aligned} u'' &= v && \text{in } (x_0, 1), \\ u(x_0, \varepsilon) &= u(1, \varepsilon) = 0, \\ \varepsilon v'' + h(u)v' - g(x, u, u')v &= 0 && \text{in } (x_0, 1), \\ v(x_0, \varepsilon) &= v^*, && v(1, \varepsilon) = v_1. \end{aligned} \tag{T2}$$

Since  $v^* = 0$ ,  $v_0 < 0$ , and  $v_1 > 0$ , applying the result of Case 1 to (T2) and applying the result of Case 2 to (T1), we have  $v(x, \varepsilon) \rightarrow 0 + BL(0)$  as  $\varepsilon \rightarrow 0^+$  in  $(0, x_0)$  and  $v(x, \varepsilon) \rightarrow 0 + BL(1)$  as  $\varepsilon \rightarrow 0^+$  in  $(x_0, 1)$ . Therefore, we conclude that

$$v(x, \varepsilon) \rightarrow 0 + BL(0) + BL(1) \quad \text{as } \varepsilon \rightarrow 0^+ \text{ in } [0, 1].$$

Case 4.  $v_0 > 0$ ,  $v_1 < 0$ . We note that this case is just the reflection of Case 3, and so we have  $v(y, \varepsilon) \rightarrow 0 + BL(0) + BL(1)$  as  $\varepsilon \rightarrow 0^+$  in  $[0, 1]$ .

Part (II).  $f(u, u') = h(u')$ . The difference between (I) and (II) is that there is no relation between the behavior of  $h(u')$  and  $h(\alpha_1)$  or  $h(\beta_1)$ , in general. However, the reduced equation,  $f(u, u')v' + g(x, u, u')v = 0$ , has only the trivial solution, and so the asymptotic behavior of the solution  $v(x, \varepsilon)$  of the coupled system (6.1) in this case is not as complicated as in Models I and II. In fact, we only need to prove that  $v(x, \varepsilon) \rightarrow 0$  as  $\varepsilon \rightarrow 0^+$  for  $v_0 \geq 0$ ,  $v_1 \geq 0$ , the three cases  $v_0 \leq 0$ ,  $v_1 \leq 0$ ;  $v_0 < 0$ ,  $v_1 > 0$ ; and  $v_0 > 0$ ,  $v_1 < 0$ , can be reduced to the first case. It is convenient to consider the subspaces  $v_0 = 0$ ,  $v_1 > 0$ ;  $v_0 > 0$ ,  $v_1 = 0$ ; and  $v_0 > 0$ ,  $v_1 > 0$  separately.

(i)  $v_0 = 0$ ,  $v_1 > 0$ . We define  $\alpha_1(x, \varepsilon) = \zeta v_1(x^2 - x)/2$ ,  $\beta_1(x, \varepsilon) = 0$ ,  $\alpha_2(x, \varepsilon) = 0$ , and  $\beta_2(x, \varepsilon) = w_R(x, \varepsilon) + \Gamma(\varepsilon)$ , where  $w_R(x, \varepsilon) = |v_1| \exp[-(g^*)^{1/2}(1-x)]$ ,  $\Gamma(\varepsilon) = \varepsilon\rho/g^*$ , and  $\zeta$  and  $\rho$  are positive constants to be determined. We note that  $\alpha_1(x, \varepsilon) \leq 0$  in  $[0, 1]$ ; it is clear that conditions (A1), (AB1), and (AB2) are satisfied. From Taylor's Theorem, we have

$$\begin{aligned} -h(u')\beta_2' + g^*\beta_2 - \varepsilon\beta_2'' &\geq g^*\beta_2 - |h(u')w_R'(x, \varepsilon)| - \varepsilon\beta_2'' \\ &\geq \varepsilon\rho - |h(u')w_R'(x, \varepsilon)| \end{aligned}$$

since  $h(u') w'_R(x, \varepsilon) > 0$  in  $[1 - \delta, 1]$  for some  $\delta > 0$ . In the rest of the interval  $[0, 1]$ ,

$$w'_R(x, \varepsilon) = |v_1| \left( \frac{g^*}{\varepsilon} \right)^{1/2} \exp \left[ - \left( \frac{g^*}{\varepsilon} \right)^{1/2} (1 - x) \right],$$

and so for sufficiently small  $\varepsilon$ , we have  $|h(u') w'_R(x, \varepsilon)| \leq c_1 \varepsilon$  for some constant  $c_1 > 0$ . Choose  $\rho \geq c_1$ , then we see that  $-h(u') \beta'_2 + g^* \beta_2 - \varepsilon \beta''_2 \geq 0$ . Thus (A3) holds. To prove (A2), since  $\beta_1(x, \varepsilon) = \alpha_2(x, \varepsilon) = 0$  and  $\alpha'_1(x, \varepsilon) = \zeta v_1$ , choose  $\zeta v_1 \geq |v_1| + \varepsilon \rho / g^*$ , then we have  $\alpha''_1 \geq \beta_2$ . Therefore,  $0 \leq v(x, \varepsilon) \leq |v_1| \exp[-(g^*/\varepsilon)^{1/2}(1-x)] + \Gamma(\varepsilon)$ . It follows that  $v(x, \varepsilon) \rightarrow 0 + BL(1)$  as  $\varepsilon \rightarrow 0^+$  in  $[0, 1]$ . If  $v_0 > 0, v_1 = 0$ . This case is the reflection of case (i). We conclude that  $v(x, \varepsilon) \rightarrow 0 + BL(0)$  as  $\varepsilon \rightarrow 0^+$  in  $[0, 1]$ .

(ii)  $v_0 > 0, v_1 > 0$ . Since both boundary values  $v_0$  and  $v_1$  are positive, we expect that there are boundary layers at both endpoints  $x = 0$  and  $x = 1$ . We define, for  $0 \leq x \leq 1$  and  $\varepsilon > 0$ ,  $\alpha_1(x, \varepsilon) = w(v_0 + v_1)(x^2 - x)/2$ ,  $\beta_1(x, \varepsilon) = 0, \alpha_2(x, \varepsilon) = 0$ , and

$$\beta_2(x, \varepsilon) = w_R(x, \varepsilon) + w_L(x, \varepsilon) + \Gamma(\varepsilon),$$

where  $\Gamma(\varepsilon) = \varepsilon \tau / g^*$  for  $\tau$  is a positive constant to be determined,  $w_R(x, \varepsilon)$  is defined in (i), and  $w_L(x, \varepsilon) = |v_0| \exp[-(g^*/\varepsilon)^{1/2}x]$ . We note that  $h(u') w'_R(x, \varepsilon) > 0$  in  $[1 - \delta, \delta]$ ,  $h(u') w'_L(x, \varepsilon) > 0$  in  $[0, \delta]$  for some  $\delta > 0$ , and

$$|h(u') \{w'_R(x, \varepsilon) + w'_L(x, \varepsilon)\}| \leq c_3 \varepsilon \quad \text{in } (\delta, 1 - \delta)$$

for  $\varepsilon$  sufficiently small and some constant  $c_3 > 0$ . If we now choose  $\tau \geq c_3$  and  $w(v_0 + v_1) \geq |v_0 + v_1| + \varepsilon \tau / g^*$ , then it follows that

$$\begin{aligned} -h(u') \beta'_2 + g^* \beta_2 - \varepsilon \beta''_2 &= -h(u')(w'_R + w'_L) + g^* \beta_2 - \varepsilon \beta''_2 \\ &\geq \varepsilon \tau - |h(u')(w'_R + w'_L)| \geq 0 \quad \text{in } [0, 1] \end{aligned}$$

and

$$\alpha''_1(x, \varepsilon) = w(v_0 + v_1) \geq \beta_2.$$

Therefore, we finish the proof to have  $0 \leq v(x, \varepsilon) \leq \beta_2(x, \varepsilon)$  in  $[0, 1]$ , that is,

$$v(x, \varepsilon) \rightarrow 0 + BL(0) + BL(1) \quad \text{as } \varepsilon \rightarrow 0^+ \quad \text{in } [0, 1].$$

We expect that this result is true for the more general system

$$\begin{aligned}
 u'' &= k(v), & \text{in } [0, 1] \\
 u(0, \varepsilon) &= u(1, \varepsilon) = 0 \\
 \varepsilon v'' + f(u, u')v' - g(x, u, u')v &= 0, & \text{in } [0, 1] \\
 v(0, \varepsilon) &= v_0, & v(1, \varepsilon) = v_1,
 \end{aligned}$$

where  $k(v)$  is a continuous function and  $\partial k/\partial v$  does not change sign for  $x$  in  $(0, 1)$ .

## 6. STABILITY OF THE STEADY-STATE SOLUTION

In the last section of this paper, we turn to examine the stability of the steady-state solution  $U = U(x, \varepsilon)$ ,  $V = V(x, \varepsilon)$  with respect to the perturbations in the initial data as  $\varepsilon \rightarrow 0^+$ . Consider the time-dependent version of (1.1)

$$\begin{aligned}
 u_{xx} - v &= u_t, & (x, t) \text{ in } (0, 1) \times (0, \infty), \\
 u(x, 0, \varepsilon) &= \phi_1(x, \varepsilon), & x \text{ in } [0, 1], \\
 u(0, t, \varepsilon) &= u(1, t, \varepsilon) = 0, & t \text{ in } [0, \infty), \\
 \varepsilon v_{xx} + f(u, u_x)v_x - g(x, u, u_x)v &= v_t, & (x, t) \text{ in } (0, 1) \times (0, \infty), \\
 v(x, 0, \varepsilon) &= \phi_2(x, \varepsilon), & x \text{ in } [0, 1], \\
 v(0, t, \varepsilon) &= v_0, & v(1, t, \varepsilon) = v_1, t \text{ in } [0, \infty),
 \end{aligned} \tag{6.1}$$

where  $f(u, u_x) = h(u)$  or  $h(u_x)$ ,  $\partial h/\partial y > 0$  for  $h = h(y)$ , and  $g(x, u, u_x) \geq 0$ . We assume that  $\|\phi_i\|_\infty$  ( $i = 1, 2$ ) is bounded in  $[0, 1] \times [0, \infty)$  as  $\varepsilon \rightarrow 0^+$ , where  $\|\psi\|_\infty$  is defined as the  $L^\infty$ -norm of  $y$ . In other words,  $(U, V)$  is the solution of the system of equations

$$\begin{aligned}
 U_{xx} &= V & \text{in } (0, 1), \\
 U(0, \varepsilon) &= U(1, \varepsilon) = 0, \\
 \varepsilon V_{xx} + f(U, U_x)V_x - g(x, U, U_x)V &= 0 & \text{in } (0, 1), \\
 V(0, \varepsilon) &= v_0, & V(1, \varepsilon) = v_1,
 \end{aligned} \tag{6.2}$$

that displays boundary layer or interior layer behavior as  $\varepsilon \rightarrow 0^+$ . We want to examine whether the solution  $(U, V)$  of (6.2) is stable or asymptotically stable with respect to perturbations of the initial data. We use the criteria

which Howes gave in [6, 7] to test the stability of  $(U, V)$  for all boundary values  $v_0$  and  $v_1$ .

Considering the  $u$ -equation  $u_t = u_{xx} - v$  of (6.1), we see from the maximum principle that  $|u| \leq \max |v|$  for all  $(x, t)$  in  $[0, 1] \times (0, \infty)$ ; in particular,  $|U| \leq \max |V|$ . Thus, it is sufficient to test the stability of the solution  $V$  of (6.2). We introduce the perturbations

$$w = w(x, t, \varepsilon) = u(x, t, \varepsilon) - U(x, \varepsilon), \quad z = z(x, t, \varepsilon) = v(x, t, \varepsilon) - V(x, \varepsilon)$$

into (6.1) and define

$$\begin{aligned} A(x, w, w_x, \varepsilon) &= f(w + U, w_x + U_x), \\ B(x, w, w_x, z, \varepsilon) &= -g(x, w + U, w_x + U_x)z \\ &\quad + [f(w + U, w_x + U_x) - f(U, U_x)]V_x \\ &\quad + [g(x, w + U, w_x + U_x) - g(x, U, U_x)]V, \\ y_1(x, \varepsilon) &= \phi_1(x, \varepsilon) - U(x, \varepsilon), \quad y_2(x, \varepsilon) = \phi_2(x, \varepsilon) - V(x, \varepsilon). \end{aligned}$$

Here  $f(w + U, w_x + U_x) = h(w + U)$  or  $h(w_x + U_x)$  and  $f(U, U_x) = h(U)$  or  $h(U_x)$ .

The perturbations are governed then by the system

$$\begin{aligned} w_{xx} &= z + w_t, \\ w(0, t, \varepsilon) &= w(1, t, \varepsilon) = 0, \quad w(x, 0, \varepsilon) = \psi_1(x, \varepsilon), \\ \varepsilon z_{xx} + A(x, w, \varepsilon)z_x + B(x, w, w_x, z, \varepsilon) &= z_t, \\ z(x, 0, \varepsilon) &= \psi_2(x, \varepsilon), \quad z(0, t, \varepsilon) = z(1, t, \varepsilon) = 0 \end{aligned} \tag{6.3}$$

We now turn to testing the stability of the zero solution of (6.3). Since  $-\partial B/\partial z = g(x, u, u_x) \geq 0$  for all  $(x, t)$  in  $[0, 1] \times [0, \infty)$ , it is clear that the zero solution of (6.3) is stable with respect to all initial perturbations as  $\varepsilon \rightarrow 0^+$ . It follows that the solution  $V$  of (6.2) is stable for all initial perturbations as  $\varepsilon \rightarrow 0^+$ . For given boundary values  $v_0$  and  $v_1$  which are not both zero, we may ask under what conditions  $V$  is asymptotically stable. To answer this question, we need to find a positive constant  $c$  such that either  $|A(x, w, w_x, \varepsilon)| \geq c$  or  $-\partial B/\partial z \geq c$ ; cf. [6, 7].

We note that  $|f(w + U, w_x + U_x)| \geq k > 0$  in  $\Delta \times (0, \infty)$ , where  $\Delta = \{[0, 1] - T\}$ ,  $T = \{x^* \mid x^* \text{ in } (0, 1) \text{ and } f(x^*, u(x^*), u'(x^*)) = 0\}$ . Therefore, if (i)  $f(u, u_x) = h(u)$ ,  $g(x, u, u_x) \equiv 0$ , for some small  $\delta$ ,  $\delta^* > 0$ , we have

- (1)  $\Delta = [\delta, 1 - \delta]$  if  $v_0 \geq 0, v_1 \geq 0$  or  $v_0 \leq 0, v_1 \leq 0$ ;
- (2)  $\Delta = [\delta, x_0 - \delta^*] \cup [x_0 + \delta^*, 1 - \delta]$  if  $v_0 > 0, v_1 < 0$  or  $v_0 < 0, v_1 > 0$ .

Then for  $\|\psi_2\|_\infty \leq \eta$ , we have  $|z(x, t, \varepsilon)| \leq C \exp(-\sigma t)$  in  $\Omega = \mathcal{A} \times (0, \infty) \times (0, \varepsilon_0]$ ,  $0 < \varepsilon_0 \leq \varepsilon$ , where  $C = \|\psi_2\|_\infty [2e^{-\lambda} - 1]$  and  $\sigma = r/(e^{-\lambda} - 1)$ , for  $\lambda = -r/k + O(\varepsilon)$  a negative root of the equation  $\varepsilon\lambda^2 + k\lambda + r = 0$ . Therefore,  $z(x, t, \varepsilon)$  is asymptotically stable in  $\Omega$ . That is, the solution  $V(x, \varepsilon)$  of (6.2) is asymptotically stable in  $\mathcal{A}$ .

(ii)  $f(u, u_x) = h(u_x)$ ,  $g(x, u, u_x) \equiv 0$ . In this case, there exist  $x_m$  in  $(0, 1)$ ,  $m = 1, 2, \dots, 6$ , such that

$$h(w_x(x_n, t, \varepsilon) + U_x(x_n, \varepsilon)) = 0 \quad \text{for } (II_n) \text{ if } n = 1, 2;$$

$$h(w_x(x_m, t, \varepsilon) + U_x(x_m, \varepsilon)) = 0 \quad \text{for } (II_n) \text{ if } n = 3, 4 \text{ and } m = n \text{ or } n + 1,$$

where

$$(II_1) := \{v_0 \geq 0, v_1 \geq 0\}, \quad (II_2) := \{v_0 \leq 0, v_1 \leq 0\},$$

$$(II_3) := \{v_0 < 0, v_1 > 0\}, \quad \text{and} \quad (II_4) := \{v_0 > 0, v_1 < 0\}.$$

Thus there exist positive constants  $k_n$  ( $n = 1, 2, 3, 4$ ) such that

$$|h(w_x + U_x)| \geq k_n > 0 \quad \text{in } (0, 1) \times (0, \infty) - \omega_n, \quad n = 1, 2, 3, 4,$$

where

$$\omega_n = I_n \times (0, \infty), \quad n = 1, 2, 3, 4,$$

$$I_n = (x_n - \delta_n, x_n + \delta_n) \quad \text{if } n = 1 \text{ or } 2,$$

$$I_n = \{(x_n - \delta_n, x_n + \delta_n) \cup (x_{n+1} - \delta_{n+1}, x_{n+1} + \delta_{n+1})\} \\ \text{if } n = 3 \text{ or } 4,$$

for some  $\delta_n$  such that  $0 < \delta_n < 1$ . Therefore, by the same arguments as in above, we conclude that the zero solution of (6.2) in this case is asymptotically stable with respect to all initial perturbations as  $\varepsilon \rightarrow 0^+$  in  $\Omega_n = \omega_n \times (0, \varepsilon_0]$  for  $(II_n)$ ,  $n = 1, 2, 3, 4$ . This is equivalent to saying that the steady-state solution  $V$  of (6.1) is asymptotically stable with respect to all initial perturbations as  $\varepsilon \rightarrow 0^+$  in  $\mathcal{A}_n = \{[\delta, 1 - \delta] - I_n\}$  for  $(II_n)$ ,  $n = 1, 2, 3, 4$ .

(iii)  $f(u, u_x) = h(u)$  or  $h(u_x)$ ,  $g(x, u, u_x) \geq g^* > 0$ . In this final case, since  $-\partial B/\partial z = g(x, u, u_x) \geq g^* > 0$ , the  $B(x, w, w_x, z, \varepsilon)$ -term dominates the stability of the solution  $z$ . We have therefore that for all initial perturbations  $\psi_2$ ,

$$|z(x, t, \varepsilon)| \leq \|\psi_2\|_\infty \exp(-g^*t) \quad \text{in } \mathcal{A} = (0, 1) \times (0, \infty) \times (0, \varepsilon_0].$$

It follows that the zero solution of (6.3) is asymptotically stable with respect to all initial perturbations as  $\varepsilon \rightarrow 0^+$ , which implies the asymptotic stability of the steady-state solution  $V$  of (6.1).

To summarize our results in this section, we conclude that the steady-state solution  $(U, V)$  of (6.1) is stable with respect to all initial perturbations as  $\varepsilon \rightarrow 0^+$  in  $[0, 1]$ . Moreover,  $V$  is asymptotically stable for  $x$  in  $[\delta, 1 - \delta]$  when  $g(x, u, u') \geq g^* > 0$  and  $V$  is asymptotically stable for  $x$  in  $\{[0, 1] - \{x^*\}\}$  when  $g(x, u, u') \equiv 0$ , where  $x^*$  in  $(0, 1)$  and  $f(x^*, u(x^*), u'(x^*)) = 0$ .

#### ACKNOWLEDGMENT

The author expresses the deepest gratitude to her advisor Professor F. A. Howes for his guidance and encouragement.

#### REFERENCES

1. K. W. CHANG AND F. A. HOWES, "Nonlinear Singular Perturbation Phenomena: Theory and Applications," Springer-Verlag, New York, 1984.
2. F. W. DORR, Some examples of singular perturbation problems with turning points, *SIAM J. Math. Anal.* **1** (1970), 141–146.
3. F. W. DORR AND S. V. PARTER, Singular perturbations of nonlinear boundary value problems with turning points, *J. Math. Anal. Appl.* **29** (1970), 273–293.
4. F. W. DORR, S. V. PARTER, AND L. F. SHAMPINE, Applications of the maximum principle to singular perturbation problems, *SIAM Rev.* **15** (1973), 43–88.
5. F. A. HOWES, "Some Old and New Results on Singularly Perturbed Boundary Value Problems" (R. E. Meyer and S. V. Parter, Eds.), pp. 41–85, Academic Press, New York, 1980.
6. F. A. HOWES, Some stability results for advection-diffusion equations, *Stud. Appl. Math.* **74** (1986), 35–53.
7. F. A. HOWES, Some stability results for advection-diffusion equations, II, *Stud. Appl. Math.* **75** (1986), 153–162.
8. F. A. HOWES AND S. SHAO, Asymptotic analysis of model problems for a coupled system, *Nonlinear Anal.* **13**, No. 9 (1989), 1013–1024.
9. W. RUDIN, "Real and Complex Analysis," McGraw-Hill, New York, 1974.
10. S. SHAO, "Asymptotic Behavior of Solutions of Model Problems for a Coupled System," Ph.D. Thesis, University of California at Davis, June 1989.